

Invariants of finite group schemes

Serge Skryabin

Chebotarev Research Institute, Kazan, Russia

Let k be an algebraically closed field, G a finite group scheme over k operating on a scheme X over k . Under assumption that X can be covered by G -invariant affine open subsets the classical results in [3] and [14] describe the quotient X/G . In case of a free action X is known to be a principal homogeneous G -space over X/G . Furthermore, the category of G -linearized quasi-coherent sheaves of $\mathcal{O}_X$ -modules is equivalent then to the category of quasi-coherent sheaves of $\mathcal{O}_{X/G}$ -modules.

In this paper we attempt to describe the situation when generic stabilizers of points on X are nontrivial. To avoid technical complications we assume that X is an algebraic variety, although the results can be extended to reduced schemes. The stabilizer G_x of a rational point $x \in X$ is a subgroup scheme of G , and we define its index $(G : G_x)$ by analogy with the ordinary finite groups. A point x is *regular* with respect to the action of G if the index $(G : G_x)$ attains the maximal possible value $q(X)$. Theorem 2.1 shows that the set $X_{G\text{-reg}}$ of all regular points is an open G -invariant subset of X , the restriction to which of the canonical morphism $\pi : X \rightarrow X/G$ is finite flat of degree $q(X)$. For every $x \in X_{G\text{-reg}}$ the fibre $\pi^{-1}(\pi(x))$ is G -equivariantly isomorphic with the quotient $G_x \backslash G$ and there is a bijective correspondence between the G -invariant closed subschemes of $X_{G\text{-reg}}$ and the closed subschemes of $\pi(X_{G\text{-reg}})$. We prove also that the field of rational functions $k(X)$ has degree $q(X)$ over the subfield of G -invariants $k(X)^G$. The arguments used in [3] and [14] are essential ingredients in our approach too. At the same time, what we prove is not quite a generalization of the classical results as we need more restrictions on X .

If $X_{G\text{-reg}} = X$ then the equivalence between categories of sheaves mentioned at the beginning extends to our settings in only as much as we restrict to G -linearized quasi-coherent sheaves generated locally by G -invariant sections (Proposition 3.2). Suppose that $\mathcal{F}$ is an arbitrary G -linearized coherent sheaf of $\mathcal{O}_X$ -modules. In Theorem 3.3 we describe an open G -invariant subset $U \subset X$ such that the sheaf of $\mathcal{O}_{X/G}$ -modules $(\pi_* \mathcal{F})^G$ is locally free of rank s over the open subset $\pi(U) \subset X/G$, where s is equal to the minimum dimension of the subspaces of G_x -invariant elements $\mathcal{F}(x)^{G_x}$ in the finite dimensional G_x -modules $\mathcal{F}(x) = \mathcal{F}_x \otimes_{\mathcal{O}_x} k(x)$, $x \in U$ (here $\mathcal{F}_x$ denotes the stalk of $\mathcal{F}$ at x and $k(x)$ the residue field of the local ring $\mathcal{O}_x$). In particular, $(\mathcal{F} \otimes_{\mathcal{O}_X} k(X))^G$ has dimension s over $k(X)^G$. We use this result to describe the G -socle of the G -module $\mathcal{F} \otimes_{\mathcal{O}_X} k(X)$ in Corollary 3.4. Given a point $x \in X_{G\text{-reg}}$ such that $\mathcal{F}_x$ is a free $\mathcal{O}_x$ -module, we show in Theorem 3.6 that $\mathcal{F}(x)$ is an injective G_x -module if and only if there exists a G -invariant affine open neighbourhood U of x such that $\mathcal{F}|_U$ is projective in the category of G -linearized sheaves of quasi-coherent $\mathcal{O}_U$ -modules. Moreover, $\mathcal{F}(U)$ and $\mathcal{F} \otimes_{\mathcal{O}_X} k(X)$ are injective G -modules in this case. To simplify the statements of results we actually consider only affine varieties X and speak about modules over the function algebra $k[X] = \mathcal{O}_X(X)$ rather than quasi-coherent sheaves. A G -linearization on a $k[X]$ -module is just a G -module structure subject to a certain compatibility requirement.

Let us call a group scheme linearly reductive if all its representations are completely reducible. Theorem 4.2 says that a point $x \in X$ has a linearly reductive stabilizer G_x if and only if x is contained in a G -invariant affine open subset $U \subset X$

such that $k[U]$ is an injective G -module. This turns out to be quite a general fact. Unlike results in previous sections X can be here any scheme over k . When X is a variety, the set of points with linearly reductive stabilizers is nonempty if and only if $k(X)$ is an injective G -module. Moreover, the structure of $k(X)$ as a G -module is completely determined in this case.

If $\text{char } k = 0$, any finite group scheme over k is constant. Then for all x in a nonempty open subset of X the stabilizer G_x coincides with the largest subgroup of G acting trivially on the whole X . This is the reason why our results present an interest mainly for fields of characteristic $p > 0$. In particular, if $G = \mathfrak{G}(\mathfrak{g})$ is the group scheme of height one corresponding to a finite dimensional p -Lie algebra $\mathfrak{g}$ then the actions of G on X correspond to actions of $\mathfrak{g}$ by derivations of the structure sheaf $\mathcal{O}_X$. Probably A. Milner was the first who observed that the degree of $k(X)$ over the subfield of $\mathfrak{g}$ -invariants $k(X)^{\mathfrak{g}}$ can be expressed in terms of Lie algebra stabilizers of points on X . He considered the special case of the adjoint representation of $\mathfrak{g}$ on its symmetric algebra $S(\mathfrak{g})$ and used the fact just mentioned to derive a lower bound for the maximum dimension of irreducible $\mathfrak{g}$ -modules [12].

In fact we are able to generalize Theorem 2.1 to the actions of not necessarily finite dimensional p -Lie algebras (Theorem 5.2). Moreover, if X is a smooth affine variety and $f_1, \dots, f_n$ are $\mathfrak{g}$ -invariant regular functions on X , taken in a suitable number, then $k[X]^{\mathfrak{g}}$ is generated by $f_1, \dots, f_n$ over the subalgebra $k[X]^{(p)}$ of p -th powers in $k[X]$ provided that the differentials $d_x f_1, \dots, d_x f_n$ are linearly independent at all points x in an open subset of X whose complement has codimension at least 2 (Theorem 5.4). In this case $k[X]^{\mathfrak{g}}$ is free over $k[X]^{(p)}$ and is a locally complete intersection ring. A similar result is valid for invariants of Frobenius kernels of reduced algebraic groups. This generalizes the work of Friedlander and Parshall [6], and Donkin [5] who considered, respectively, the adjoint and the conjugating actions of a semisimple algebraic group. We discuss yet another example of the adjoint action of the Jacobson-Witt algebra W_n . Other applications to invariants of Lie algebras of Cartan type will be a subject of separate papers.

I would like to thank the referee for making comments and correction in attributing the formula for the p -th powers of derivations in section 5.

1. Preliminaries.

Let k be an algebraically closed field. It is the ground field for our considerations, so that the functors $\otimes$, Hom etc. are assumed to be taken over k unless the base ring is indicated explicitly. Let G be a finite group scheme over k and $k[G]$ the associated finite dimensional Hopf algebra. We will be considering a group action $\mu : X \times G \rightarrow X$ of G on a scheme X over k from the right. By [3] a scheme can be regarded as a functor on the category of commutative k -algebras. For each commutative algebra K the group $G(K)$ operates on $X(K)$, and this action is natural in K . If X is affine with algebra $k[X]$ then the quotient X/G is defined to be $\text{Spec } k[X]^G$ where $k[X]^G \subset k[X]$ is the subalgebra of G -invariants. More generally, if X can be covered by G -invariant affine open subsets U , then X/G is obtained by patching together the affine quotients U/G . We list below the properties of the canonical morphism $\pi : X \rightarrow X/G$ assuming X to be of finite type (see [3], Ch. III, §2, 6.1 and [14], Ch. III, §12):

- (1) π is finite and surjective,
- (2) the set-theoretic fibers of π coincide with the orbits of the group $G(k)$,
- (3) X/G has the quotient topology with respect to π ; in particular, π is open,

(4) if $U \subset X$ is a G -invariant open subscheme, then $U/G \cong \pi(U)$ and $U = \pi^{-1}(\pi(U))$.

According to [3], Ch. III, §2, 2.3 the action is said to be free if $G(K)$ operates freely on $X(K)$ for each commutative algebra K . In case of a free action π is finite flat of degree $|G| = \dim k[G]$ (which means that $k[U]$ are free of rank $|G|$ over $k[U]^G$ for a suitable covering of X by G -invariant affine open subsets) and the canonical morphism $\nu = (p_1, \mu) : X \times G \rightarrow X \times_{X/G} X$, where $p_1 : X \times G \rightarrow X$ denotes the projection, is an isomorphism.

If $G' \subset G$ is a subgroup scheme, we let $G' \backslash G$ denote the quotient with respect to the action of G' on G by left translations. Then $G' \backslash G$ is a finite scheme with the algebra $k[G' \backslash G] \cong k[G]^{G'}$, the invariants with respect to the left regular representation of G' on $k[G]$. Define the *index* of G' in G to be $(G : G') = \dim k[G' \backslash G]$. The algebra $k[G]$ is a free module of rank $|G'|$ over $k[G' \backslash G]$. Hence $|G| = (G : G') \cdot |G'|$. The index can be interpreted in terms of dual Hopf algebras. By [16] or [15] $k[G]^*$ is free both as a left and a right module over its subalgebra $k[G']^*$. Clearly, the ranks of these modules are equal to $(G : G')$.

If $x \in X(k)$ then its stabilizer $G_x \subset G$ is a subgroup scheme such that $G_x(K) = \{g \in G(K) \mid x_K g = x_K\}$ for each commutative algebra K , where x_K denotes the image of x in $X(K)$. Let $i_x : \text{Spec } k \rightarrow X$ be the morphism corresponding to x and

$$\mu_x : G \cong \text{Spec } k \times G \xrightarrow{i_x \times \text{id}_G} X \times G \xrightarrow{\mu} X.$$

the orbit morphism. Then G_x coincides with the fiber of μ_x above x and μ_x factors through a morphism $\rho : G_x \backslash G \rightarrow X$. By [3], Ch. III, §3, 5.2 ρ is an immersion. In fact ρ is a closed immersion because $G' \backslash G$ has only rational points. Since the case of finite group schemes is especially easy, below we sketch a proof of an equivalent assertion for the reader's convenience:

PROPOSITION 1.1. *Suppose that $A \subset k[G]$ is a subalgebra, stable under the right regular representation of G , and $\mathfrak{m}_A = \mathfrak{m} \cap A$ where $\mathfrak{m}$ is the augmentation ideal of $k[G]$. Then $A = k[G' \backslash G]$ where G' is the stabilizer of $\mathfrak{m}_A$ in G .*

Proof. Put $B = k[G' \backslash G]$, $R = k[G]$. The group scheme G operates from the right on $X = \text{Spec } A$ and the inclusion $A \subset R$ corresponds to a G -equivariant morphism $G \rightarrow X$. The latter is the orbit morphism μ_x of the point $x \in X(k)$ corresponding to $\mathfrak{m}_A$. Since $G' = G_x$, the orbit morphism factors through $G' \backslash G$, which means that $A \subset B$. Let $K = R/R\mathfrak{m}_A$, and let $g \in G(K)$ be the point corresponding to the canonical homomorphism $R \rightarrow K$. Since the composite homomorphism $A \rightarrow R \rightarrow K$ factors through $A/\mathfrak{m}_A$, we have $x_K g = \mu_x(g) = x_K$, which yields $g \in G'(K)$. It follows that the composite $B \rightarrow R \rightarrow K$ factors through $B/\mathfrak{m}_B$ where $\mathfrak{m}_B = \mathfrak{m} \cap B$. In other words, $\mathfrak{m}_B = R\mathfrak{m}_A \cap B$. However, $R\mathfrak{m}_A \cap B = B\mathfrak{m}_A$ because R is free over B . Hence $B = k + \mathfrak{m}_B = A + B\mathfrak{m}_A$. Since R is finite over A , the map $G(k) \rightarrow X(k)$ determined by μ_x is surjective. This means that $G(k)$ transitively permutes the maximal ideals of A . Then $B = A + B\mathfrak{n}$ for all maximal ideals $\mathfrak{n}$ of A . An application of Nakayama's lemma yields $B = A$. $\square$

Suppose now that A is any unital associative algebra on which G operates by automorphisms. This means that A has a G -module structure and for each commutative algebra K the group $G(K)$ operates on $A \otimes K$ via automorphisms. We call A a G -algebra in this case. By an (A, G) -module we mean a right A -module M equipped with an additional G -module structure such that the A -module structure

map $M \otimes A \rightarrow M$ is G -equivariant. Denote by $\mathcal{M}_A$ the category of (A, G) -modules. The morphisms in $\mathcal{M}_A$ are maps which are simultaneously A -module and G -module homomorphisms.

This definition is meaningful for an arbitrary group scheme. When G is finite, the category of G -modules is equivalent to the category of left $k[G]^*$ -modules (see [9], Part I, Ch. 8), and $\mathcal{M}_A$ is equivalent to the category of left modules over the smash product algebra $A^{\text{op}} \# k[G]^*$ where A^{op} is the algebra A with the opposite multiplication. When A is commutative, $A^{\text{op}} = A$. We refer the reader to [19] or [13] concerning the precise definition of smash products. Here we just point out that $A \# k[G]^*$ contains A and $k[G]^*$ as subalgebras and the multiplication map $A \otimes k[G]^* \rightarrow A \# k[G]^*$ is bijective. One of our tools is the following theorem whose interpretations can be found in [2], [10], [20]:

IMPRIMITIVITY THEOREM. *Suppose that $A = k[G' \setminus G]$, and let $\mathfrak{m}_A$ be the augmentation ideal of A . Then the functor $M \mapsto M/M\mathfrak{m}_A$ is an equivalence between $\mathcal{M}_A$ and the category of G' -modules. If $M \in \mathcal{M}_A$, then there is an isomorphism of G -modules $M \cong \text{ind}_{G'}^G M/M\mathfrak{m}_A$.*

Given a G' -module V , the induced G -module is $\text{ind}_{G'}^G V = (V \otimes k[G])^{G'}$. To be precise, we compute the G' -invariants with respect to the tensor product of the G' -module structure on V and the left regular G' -module structure on $k[G]$, and we use the right regular G -module structure on $k[G]$ to get one on $\text{ind}_{G'}^G V$. This differs from the conventions adopted in [9]. In terms of the dual algebras

$$\text{ind}_{G'}^G V \cong \text{Hom}_{k[G']^*}(k[G]^*, V).$$

It follows that $\dim \text{ind}_{G'}^G V = (G : G') \cdot \dim V$. We mention also that the injective G -modules are projective, and vice versa, because finite dimensional Hopf algebras are Frobenius (see [19]). Waterhouse proved that every finite group scheme is geometrically reductive [22].

Below we prove two lemmas. Suppose that A is a commutative integral domain and K its field of fractions. For a prime ideal $\mathfrak{p}$ of A denote $k(\mathfrak{p}) = A_{\mathfrak{p}}/A_{\mathfrak{p}}\mathfrak{p}$.

LEMMA 1.2. *Let F be a finitely generated projective A -module, F' its submodule. Denote by $I(\mathfrak{p})$ the image of the canonical map $F' \otimes_A k(\mathfrak{p}) \rightarrow F \otimes_A k(\mathfrak{p})$ and put $q = \dim_K F' \otimes_A K$. Then:*

- (1) $q = \max_{\mathfrak{p} \in \text{Spec } A} \dim_{k(\mathfrak{p})} I(\mathfrak{p})$.
- (2) *The subset $U = \{\mathfrak{p} \in \text{Spec } A \mid \dim_{k(\mathfrak{p})} I(\mathfrak{p}) = q\}$ is Zariski open and coincides with the subset $U' = \{\mathfrak{p} \in \text{Spec } A \mid F'_{\mathfrak{p}} \text{ is a direct summand of } F_{\mathfrak{p}}\}$.*
- (3) *If $U = \text{Spec } A$ then F' is projective of rank q and is a direct summand of F .*

Proof. We identify all localizations of F , as well as localizations of F' , with their images in $F \otimes_A K$. Let $\mathfrak{p} \in \text{Spec } A$. Suppose that $u_1, \dots, u_r \in F'_{\mathfrak{p}}$ are elements whose images in $F_{\mathfrak{p}}/\mathfrak{p}F_{\mathfrak{p}} \cong F \otimes_A k(\mathfrak{p})$ are linearly independent over $k(\mathfrak{p})$. Then $S = A_{\mathfrak{p}}u_1 + \dots + A_{\mathfrak{p}}u_r$ is a direct summand of the free $A_{\mathfrak{p}}$ -module $F_{\mathfrak{p}}$ (see [1], Ch. II, §3, Cor. 1 to Prop. 5). In particular, S is free over $A_{\mathfrak{p}}$ with a basis $u_1, \dots, u_r$. We get $r = \dim_K S \otimes_A K \leq q$. It follows $\dim_{k(\mathfrak{p})} I(\mathfrak{p}) \leq q$. By definition of q the equality holds here for $\mathfrak{p} = (0)$.

Suppose that $\mathfrak{p} \in U$. Then we can take q elements $u_1, \dots, u_q$ above, so that $r = q$. By modularity law S is a direct summand of $F'_{\mathfrak{p}}$. So $F'_{\mathfrak{p}} = S \oplus C$ where $C \subset F_{\mathfrak{p}}$ is an $A_{\mathfrak{p}}$ -submodule. Tensoring with K and comparing dimensions over K ,

we deduce that $C \otimes_A K = 0$. However, $F_{\mathfrak{p}}$ is torsion-free since $A_{\mathfrak{p}}$ is a domain. It follows $C = 0$, i.e., $F'_{\mathfrak{p}} = S$. Thus $\mathfrak{p} \in U'$.

Conversely, suppose that $\mathfrak{p} \in U'$. Then $F'_{\mathfrak{p}}$ is free of rank q over $A_{\mathfrak{p}}$ and the map $F' \otimes_A k(\mathfrak{p}) \rightarrow F \otimes_A k(\mathfrak{p})$ is injective. It follows $\dim_{k(\mathfrak{p})} I(\mathfrak{p}) = q$. There exists an $A_{\mathfrak{p}}$ -module epimorphism $\varphi : F_{\mathfrak{p}} \rightarrow F'_{\mathfrak{p}}$ which is identity on $F'_{\mathfrak{p}}$. Since F is finitely generated, $\varphi(F) \subset F'_s$ for a suitable $s \in A \setminus \mathfrak{p}$. Then $\varphi(F_s) = F'_s$, and so F'_s is a direct summand of F_s . This shows that U' is open in $\text{Spec } A$. Finally, (3) is obtained by an application of [1], Ch. II, §5, Th. 1 and §3, Cor. 1 to Prop. 12. $\square$

LEMMA 1.3. *Let F be a finitely generated projective A -module, F' and F'' its direct summands. Denote by $I'(\mathfrak{p})$, $I''(\mathfrak{p})$ the images, respectively, of $F' \otimes_A k(\mathfrak{p})$, $F'' \otimes_A k(\mathfrak{p})$ in $F \otimes_A k(\mathfrak{p})$, and put $s = \min_{\mathfrak{p} \in \text{Spec } A} \dim_{k(\mathfrak{p})} I'(\mathfrak{p}) \cap I''(\mathfrak{p})$. Then:*

(1) *The subset $U = \{\mathfrak{p} \in \text{Spec } A \mid \dim_{k(\mathfrak{p})} I'(\mathfrak{p}) \cap I''(\mathfrak{p}) = s\}$ is open in $\text{Spec } A$ and consists precisely of those $\mathfrak{p}$ for which $F'_{\mathfrak{p}} + F''_{\mathfrak{p}}$ is a direct summand of $F_{\mathfrak{p}}$.*

(2) *For all $\mathfrak{p} \in U$ the canonical map $(F' \cap F'') \otimes_A k(\mathfrak{p}) \rightarrow F \otimes_A k(\mathfrak{p})$ is an isomorphism onto $I'(\mathfrak{p}) \cap I''(\mathfrak{p})$.*

(3) *If $U = \text{Spec } A$ then the A -module $F' \cap F''$ is projective of rank s and is a direct summand of F .*

Proof. Since F' , F'' are direct summands of F , the dimensions of vector spaces $I'(\mathfrak{p})$, $I''(\mathfrak{p})$ do not depend on $\mathfrak{p}$. Then U is the set of those $\mathfrak{p} \in \text{Spec } A$ for which $I'(\mathfrak{p}) + I''(\mathfrak{p})$ has maximal possible dimension. Apply now Lemma 1.2 taking $F' + F''$ instead of F' in it. We get assertion (1) of Lemma 1.3. If $\mathfrak{p} \in U$ then $F'_{\mathfrak{p}} + F''_{\mathfrak{p}}$ is a free $A_{\mathfrak{p}}$ -module. Since $F'_{\mathfrak{p}}$ is a direct summand of $F'_{\mathfrak{p}} + F''_{\mathfrak{p}}$, the $A_{\mathfrak{p}}$ -module

$$F'_{\mathfrak{p}} / (F'_{\mathfrak{p}} \cap F''_{\mathfrak{p}}) \cong (F'_{\mathfrak{p}} + F''_{\mathfrak{p}}) / F''_{\mathfrak{p}} \quad (*)$$

is free too. Then $F'_{\mathfrak{p}} \cap F''_{\mathfrak{p}}$ is a direct summand of $F'_{\mathfrak{p}}$, hence also a direct summand of $F_{\mathfrak{p}}$. In particular, $(F' \cap F'')_{\mathfrak{p}} \cong F'_{\mathfrak{p}} \cap F''_{\mathfrak{p}}$ is free over $A_{\mathfrak{p}}$ and the map in (2) is injective. Denoting by $I(\mathfrak{p})$ the image of that map and tensoring (*) with $k(\mathfrak{p})$, we obtain an isomorphism $I'(\mathfrak{p}) / I(\mathfrak{p}) \cong (I'(\mathfrak{p}) + I''(\mathfrak{p})) / I''(\mathfrak{p})$. It follows $I(\mathfrak{p}) = I'(\mathfrak{p}) \cap I''(\mathfrak{p})$. The last assertion is a special case of Lemma 1.2(3). $\square$

2. The set of G -regular points and the properties of the quotient.

Let G be a finite group scheme operating from the right on an irreducible algebraic variety X . Suppose that X can be covered by G -invariant affine open subsets, so that X/G exists (as is well known it suffices to require that the $G(k)$ -orbit of each closed point of X is contained in an affine open subset). We will be considering only closed points of X , so that $x \in X$ means $x \in X(k)$. If $U \subset X$ is an open subset, stable under all automorphisms of X determined by the elements of $G(k)$, then the composite morphism $U \times G \rightarrow X \times G \xrightarrow{\mu} X$ factors through U , i.e., U is G -invariant. In particular, if $U \subset X$ is any open subset, $\cap_{g \in G(k)} Ug$ is a G -invariant open subset contained in U . It follows that the field of rational functions $k(X)$ is a direct limit of the G -algebras $k[U]$ where U runs through the G -invariant affine open subvarieties of X . Hence G operates on $k(X)$ by automorphisms. Put

$$q(X) = \max_{x \in X} (G : G_x),$$

$$X_{G\text{-reg}} = \{x \in X \mid (G : G_x) = q(X)\}.$$

THEOREM 2.1. $X_{G\text{-reg}}$ is a G -invariant open subset of X . Furthermore:

- (1) $\pi|_{X_{G\text{-reg}}} : X_{G\text{-reg}} \rightarrow \pi(X_{G\text{-reg}})$ is a finite flat morphism of degree $q(X)$.
- (2) For every $x \in X_{G\text{-reg}}$ the fibre of π above $\pi(x)$ is G -equivariantly isomorphic with $G_x \backslash G$.
- (3) The G -invariant closed subschemes Z of $X_{G\text{-reg}}$ are in a bijective correspondence with the closed subschemes W of $\pi(X_{G\text{-reg}})$. If Z and W correspond to each other, then $W \cong Z/G$ and $Z \cong W \times_{X/G} X$.
- (4) One has $(Y \times_{X/G} X)/G \cong Y$ for any scheme Y on which G operates trivially and a morphism $Y \rightarrow \pi(X_{G\text{-reg}})$.
- (5) $[k(X) : k(X)^G] = q(X)$.

We first reformulate the assertions of the theorem in the affine case.

PROPOSITION 2.2. Suppose that $X \cong \text{Spec } A$ and $X_{G\text{-reg}} = X$. Then:

- (1) A is a projective A^G -module of rank $q(X)$.
- (2) If $\mathfrak{m}$ is a maximal ideal of A and $\mathfrak{n} = \mathfrak{m} \cap A^G$ then $\mathfrak{n}A$ is a maximal G -invariant ideal of A and the algebra $A/\mathfrak{n}A$ is G -equivariantly isomorphic with $k[G_{\mathfrak{m}} \backslash G]$ where $G_{\mathfrak{m}}$ is the stabilizer of $\mathfrak{m}$ in G .
- (3) The assignment $I \mapsto I^G$ establishes a bijection between the G -invariant ideals of A and the ideals of A^G . The inverse correspondence is given by $J \mapsto JA$. The canonical maps $A^G \rightarrow (A/I)^G$ are surjective.
- (4) If B is an A^G -algebra on which G operates trivially then $(B \otimes_{A^G} A)^G \cong B$.

Proof. Given any covering of X by G -invariant open subvarieties, it suffices to prove the theorem for the induced action of G on each of these subvarieties. In particular, we may assume X to be affine. Let $A = k[X]$, $R = k[G]$, and let $\mu^* : A \rightarrow A \otimes R$, $i_x^* : A \rightarrow k$, $\mu_x^* : A \rightarrow R$ be the comorphisms of μ, i_x, μ_x , respectively. Denote by $\mathfrak{m}_x$ the maximal ideal of A consisting of functions vanishing at x .

Consider $F = A \otimes R$ as an A -module by means of the algebra homomorphism $p_1^* : A \rightarrow A \otimes R$, $a \mapsto a \otimes 1$. Clearly F is free of finite rank over A . Put

$$F' = (A \otimes 1) \cdot \mu^*(A) \subset F.$$

Then F' is an A -submodule of F generated by $\mu^*(A)$. Denote by $I(x)$ the image of the canonical map $F'/\mathfrak{m}_x F' \rightarrow F/\mathfrak{m}_x F$. We have $F/\mathfrak{m}_x F \cong A/\mathfrak{m}_x \otimes R \cong R$ and $I(x) = \mu_x^*(A)$ since $\mu_x^* = (i_x^* \otimes \text{id}_R) \circ \mu^*$. Now μ_x^* is a G -equivariant algebra homomorphism. Since G_x coincides with the stabilizer of $\mathfrak{m}_x$ in G , Proposition 1.1 ensures $\mu_x^*(A) = k[G_x \backslash G]$. Hence $\dim I(x) = (G : G_x)$. It follows that $X_{G\text{-reg}}$ coincides with the set U of those points $x \in X$ for which $\dim I(x)$ attains its maximal value $q = q(X)$. We can now apply Lemma 1.2. By (2) of the lemma U is open. Each $g \in G(k)$ determines an inner automorphism of G which induces an isomorphism $G_x \cong G_{xg}$. Hence $(G : G_{xg}) = (G : G_x)$. It follows that U is G -invariant.

Let $y \in U$ and let $O \subset X$ be the $G(k)$ -orbit of y . Then $\dim I(z) = q$ for all $z \in O$. Since O is finite, we can find $a_1, \dots, a_q \in A$ such that $\mu_z^*(a_1), \dots, \mu_z^*(a_q)$ are a basis of $I(z)$ for each $z \in O$. Furthermore, we may assume $a_1 = 1$ since $\mu_x^*(1) = 1$ for all x . Applying Lemma 1.2 to the A -submodule of F generated by $\mu^*(a_1), \dots, \mu^*(a_q)$, we see that the set U_1 of those $x \in X$ for which $\mu_x^*(a_1), \dots, \mu_x^*(a_q)$ are linearly

independent is open in X . Clearly $U_1 \subset U$. Since π is a finite morphism, the set $W = \pi(X \setminus U_1)$ is closed in X/G . Since $\pi^{-1}(\pi(y)) = O \subset U_1$, we have $\pi(y) \notin W$. Let V be an open affine neighbourhood of $\pi(y)$ in X/G . Then $\pi^{-1}(V)$ is an open affine G -invariant neighbourhood of y in X and $\pi^{-1}(V) \subset U_1$.

To prove the remainder of the theorem we can again use the local character of the assertions and pass to the actions of G on the invariant open subsets of the form $\pi^{-1}(V)$ constructed above. We may thus assume that $U_1 = X$.

By Lemma 1.2 F' is a direct summand of the A -module F and for each maximal ideal $\mathfrak{m}$ of A the localization $F'_{\mathfrak{m}}$ is free of rank q over $A_{\mathfrak{m}}$. If $\varphi : A^q \rightarrow F'$ is the A -module homomorphism sending the standard generators of A^q to $\mu^*(a_1), \dots, \mu^*(a_q)$ then the localizations of φ at maximal ideals of A are all isomorphisms. Hence φ is itself an isomorphism, i.e., F' is a free A -module with a basis $\mu^*(a_1), \dots, \mu^*(a_q)$. Hence for each $a \in A$ there are $b_1, \dots, b_q \in A$ such that

$$\mu^*(a) = \sum (b_i \otimes 1) \cdot \mu^*(a_i). \quad (*)$$

Let $\varepsilon : R \rightarrow k$ be the counit and $m^* : R \rightarrow R \otimes R$ the comultiplication maps. Applying $\text{id}_A \otimes \varepsilon$ to both sides of $(*)$, we get $a = \sum b_i a_i$ since $(\text{id}_A \otimes \varepsilon) \circ \mu^* = \text{id}_A$. Applying $\mu^* \otimes \text{id}_R$ and $\text{id}_A \otimes m^*$ to both sides of $(*)$, and taking into account the identity $(\mu^* \otimes \text{id}_R) \circ \mu^* = (\text{id}_A \otimes m^*) \circ \mu^*$, we get

$$\sum (\mu^*(b_i) \otimes 1) \cdot (\mu^* \otimes \text{id}_R) \mu^*(a_i) = \sum (b_i \otimes 1 \otimes 1) \cdot (\mu^* \otimes \text{id}_R) \mu^*(a_i) \quad (**)$$

in $A \otimes R \otimes R$. If $\gamma : A \rightarrow A'$ is a homomorphism of commutative algebras then $A' \otimes_A F'$ is a free A' -module with basis elements $1 \otimes \mu^*(a_i)$, $i = 1, \dots, q$. Since the canonical map $A' \otimes_A F' \rightarrow A' \otimes_A F \cong A' \otimes R$ is injective, the elements

$$(\gamma \otimes \text{id}_R) \mu^*(a_i) \in A' \otimes R, \quad i = 1, \dots, q,$$

are linearly independent over A' . Taking $A' = A \otimes R$ and $\gamma = \mu^*$, we deduce from $(**)$ that $\mu^*(b_i) = b_i \otimes 1$, that is, $b_i \in A^G$ for all i . Hence $A = A^G a_1 + \dots + A^G a_q$. If now $\sum c_i a_i = 0$ for some $c_1, \dots, c_q \in A^G$ then $\sum (c_i \otimes 1) \mu^*(a_i) = \mu^*(\sum c_i a_i) = 0$, whence $c_i = 0$ for all i . Thus A is free of rank q over A^G .

Suppose that N is an A^G -module and $M = N \otimes_{A^G} A$ is given a G -module structure by means of the comodule structure map

$$\text{id}_N \otimes \mu^* : N \otimes_{A^G} A \rightarrow N \otimes_{A^G} (A \otimes R) \cong (N \otimes_{A^G} A) \otimes R.$$

We claim that the assignment $n \mapsto n \otimes 1$ yields an isomorphism $N \cong M^G$. This amounts to showing that the exactness of the sequence of A^G -modules

$$0 \longrightarrow A^G \longrightarrow A \xrightarrow{\mu^* - p_1^*} A \otimes R$$

is preserved under tensoring with N over A^G . Now $A = A^G \oplus (A^G a_2 + \dots + A^G a_q)$ as we assume $a_1 = 1$. Next, $(\mu^* - p_1^*)(A)$ is an A^G -submodule of F generated by the elements $\mu^*(a_i) - a_i \otimes 1$, $i = 2, \dots, q$. Note that these elements together with $\mu^*(a_1) = 1 \otimes 1$ give a basis for F' over A . Then the A -submodule generated by $\mu^*(a_i) - a_i \otimes 1$, $i = 2, \dots, q$, is a direct summand of F' , hence also of F . We have seen that A^G is a direct summand of A , hence also $(\mu^* - p_1^*)(A)$ is a direct summand of F as A^G -modules. Our claim follows.

Assertion (4) of the theorem is local on Y , hence it suffices to consider an affine scheme $Y \cong \operatorname{Spec} B$. This is then a special case of what we have just proved. Next, taking $N = J$ where J is an ideal of A^G , we get $JA \cap A^G = J$ since $J \otimes_{A^G} A \cong JA$ by projectivity of A over A^G . Taking $N = A^G/J$, we deduce that the canonical map $A^G \rightarrow (A/JA)^G$ is surjective. Suppose that I is a G -invariant ideal of A . Then $\mu^*(I) \subset I \otimes R = IF$. Since on the other hand $\mu^*(A) \subset F'$ and F' is a direct summand of F , we get $\mu^*(I) \subset F' \cap IF = IF'$. Given $a \in I$, we can write therefore the expression $(*)$ with $b_i \in I$. As we have seen, this implies $a = \sum b_i a_i$ and $b_i \in A^G$. Thus $I = I^G A$, which completes the proof of (3). If $\mathfrak{n}$ is a maximal ideal of A^G then $\mathfrak{n}A$ is a maximal G -invariant ideal of A by (3). If now $\mathfrak{n} = \mathfrak{m}_x \cap A^G$ then $\mu^*(\mathfrak{n}A) = (\mathfrak{n} \otimes 1) \cdot \mu^*(A) \subset \mathfrak{m}_x \otimes R$, whence $\mu_x^*(\mathfrak{n}A) = 0$. It follows that $\ker \mu_x^* = \mathfrak{n}A$, and $A/\mathfrak{n}A \cong \mu_x^*(A)$. This proves (2). Assertion (5) follows from (1) since $k(X)^G$ is the field of fractions of the ring A^G . $\square$

COROLLARY 2.3. *If x is a smooth point of $X_{G\text{-reg}}$ then $\pi(x)$ is a smooth point of X/G .*

Proof. The local ring $\mathcal{O}_{x,X}$ is a flat extension of $\mathcal{O}_{\pi(x),X/G}$. Since $\mathcal{O}_{x,X}$ is a regular local ring, so is $\mathcal{O}_{\pi(x),X/G}$ too by [11], (21.D). $\square$

Remark. Theorem 2.1 can be generalized to the case when k is any field and X is a reduced scheme over k . In general the stabilizer G_x is a subgroup scheme of $G \otimes k(x)$ where $k(x)$ is the residue field of a point $x \in X$, and one can define $X_{G\text{-reg}}$ to be the set of all points where the function $x \mapsto (G \otimes k(x) : G_x)$ is locally constant. It can also be proved that the morphism $\nu : X_{G\text{-reg}} \times G \rightarrow X_{G\text{-reg}} \times_{\pi(X_{G\text{-reg}})} X_{G\text{-reg}}$ is finite flat. The assumption that X is reduced is needed to ensure that F' is a direct summand of F in the proof of theorem 2.1. Simple examples show what can happen without this assumption. Suppose, for instance, that $X = \operatorname{Spec} A$ where $A = k[x]$, $x^3 = 0$, and G is the cyclic order 2 group with a generator σ which acts on A as the automorphism sending x to $-x$. Here $A^G = k + kx^2$ (provided $\operatorname{char} k \neq 2$). Clearly A is not free over A^G . At the same time X contains a single point, so that $X_{G\text{-reg}} = X$.

3. G -linearized modules.

We keep our assumptions on G and X from section 2. Moreover, we assume here that X is affine. Let $R = k[G]$, $A = k[X]$ and $K = k(X)$. Recall that $\mathcal{M}_A$ denotes the category of (A, G) -modules. Denote by $\mu^M : M \rightarrow M \otimes R$ the map that gives $M \in \mathcal{M}_A$ the R -comodule structure corresponding to the G -module structure. In particular, A is an R -comodule via the map $\mu^* : A \rightarrow A \otimes R$ which is the comorphism of μ . We may view $M \otimes R$ as a module over $A \otimes R$ in a natural way. The compatibility of A - and G -module structures on M can be expressed in terms of the identity $\mu^M(ma) = \mu^M(m) \cdot \mu^*(a)$ where $m \in M$ and $a \in A$. We note also that $(S^{-1}M)^G \cong S^{-1}M^G$ for every multiplicatively closed subset $S \subset A^G$.

LEMMA 3.1. *Suppose that $X = X_{G\text{-reg}}$ and $M \in \mathcal{M}_A$. Then*

$$\begin{aligned} M^G A &= \{m \in M \mid \mu^M(m) \in (M \otimes 1) \cdot \mu^*(A)\} \\ &= \{m \in M \mid m \otimes 1 \in \mu^M(M) \cdot (A \otimes 1)\}. \end{aligned}$$

Proof. Let $F = A \otimes R$ and $F' = (A \otimes 1) \cdot \mu^*(A)$ as in the proof of Theorem 2.1. For every A -module N we consider $F_N = N \otimes R$ as an F -module in a natural

way and as an A -module by means of the homomorphism $p_1^* : A \rightarrow F$. Then $F_N \cong N \otimes_A F$. Put $F'_N = (N \otimes 1) \cdot \mu^*(A) \subset F_N$. Since F' is a direct summand of F , the canonical map $N \otimes_A F' \rightarrow N \otimes_A F$ is a split monomorphism of A -modules. Clearly its image coincides with F'_N . Hence $F'_N \cong N \otimes_A F'$.

Suppose that $m \in M$ and $\mu^M(m) \in F'_M$. To prove that $m \in M^G A$ it suffices to show that for every $x \in X$ there exists $f \in A^G$ such that $f(x) \neq 0$ and $mf \in M^G A$. Passing to suitable localizations A_f and M_f , we may thus assume as in the proof of Theorem 2.1 that F' is a free A -module with basis elements $\mu^*(a_1), \dots, \mu^*(a_q)$. Then $\mu^M(m) = \sum (m_i \otimes 1) \cdot \mu^*(a_i)$ for some $m_1, \dots, m_q \in M$. Let ε and m^* be the counit and the comultiplication in R . Applying $\text{id}_M \otimes \varepsilon$ to both sides of the equality, we get $m = \sum m_i a_i$. Applying $\mu^M \otimes \text{id}_R$ and $\text{id}_M \otimes m^*$, we get

$$\sum (\mu^M(m_i) \otimes 1) \cdot (\mu^* \otimes \text{id}_R) \mu^*(a_i) = \sum (m_i \otimes 1 \otimes 1) \cdot (\mu^* \otimes \text{id}_R) \mu^*(a_i)$$

in $M \otimes R \otimes R$. If $n_1, \dots, n_q$ are elements of an A -module N with the property that $\sum (n_i \otimes 1) \cdot \mu^*(a_i) = 0$ in F_N then $\sum n_i \otimes \mu^*(a_i) = 0$ in $N \otimes_A F'$ by the discussion at the beginning of the proof, whence $n_i = 0$ for all i . Now take $N = M \otimes R$ with the A -module structure given by means of the algebra homomorphism $\mu^* : A \rightarrow A \otimes R$. Then $A \otimes R$ operates in $N \otimes R \cong M \otimes R \otimes R$ by means of the algebra homomorphism $\mu^* \otimes \text{id}_R$, and it follows from the displayed equation above that $\mu^M(m_i) = m_i \otimes 1$, i.e., $m_i \in M^G$ for all i . Thus $m \in M^G A$.

Suppose now that $m \in M$ and $m \otimes 1 = \sum \mu^M(m_i) \cdot (b_i \otimes 1)$ for some elements $m_1, \dots, m_r \in M$ and $b_1, \dots, b_r \in A$. If $\beta : R \rightarrow B$ is the algebra homomorphism corresponding to a point $g \in G(B)$ where B is a commutative algebra then, applying $\text{id}_M \otimes \beta$ to both sides of the equality, we get $m \otimes 1 = \sum g(m_i \otimes 1) \cdot (b_i \otimes 1)$ in $M \otimes B$ (regarded as a module over $A \otimes B$). Replacing here g with g^{-1} and applying g to both sides of the equality obtained, we get $g(m \otimes 1) = \sum (m_i \otimes 1) \cdot g(b_i \otimes 1)$. If now $B = R$ and $g \in G(R)$ is the point corresponding to the identity homomorphism $R \rightarrow R$, this can be rewritten as $\mu^M(m) = \sum (m_i \otimes 1) \cdot \mu^*(b_i)$. Thus we have come to the case already considered. $\square$

PROPOSITION 3.2. *Suppose that $X = X_{G\text{-reg}}$ and denote by $\mathcal{M}'_A$ the full subcategory of $\mathcal{M}_A$ consisting of (A, G) -modules M such that $M = M^G A$. Then:*

- (1) $\mathcal{M}'_A$ is closed under taking submodules and factor modules.
- (2) The functor $M \mapsto M^G$ is an equivalence between $\mathcal{M}'_A$ and the category of A^G -modules. The inverse functor is $N \mapsto N \otimes_{A^G} A$.
- (3) If $M \in \mathcal{M}'_A$ is projective of rank r as an A -module then M^G is projective of rank r as an A^G -module.

Proof. (1) We use the same notations as in the preceding lemma. Clearly, $F_N/F'_N \cong N \otimes_A F/F'$ for every A -module N . Let N be an (A, G) -submodule of $M \in \mathcal{M}'_A$. Since F/F' is a projective A -module, the canonical map $N \otimes_A F/F' \rightarrow M \otimes_A F/F'$ is injective. It follows from the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & F'_N & \longrightarrow & F_N & \longrightarrow & N \otimes_A F/F' \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & F'_M & \longrightarrow & F_M & \longrightarrow & M \otimes_A F/F' \longrightarrow 0 \end{array}$$

that $F'_N = F_N \cap F'_M$. Now $\mu^M(N) \subset F_N$ and $\mu^M(M) \subset F'_M$, whence $\mu^M(N) \subset F'_N$. Hence $N \in \mathcal{M}'_A$ by Lemma 3.1. The assertion about factor modules is obvious.

(2) If N is an A^G -module and $M = N \otimes_{A^G} A$ then $N \cong M^G$ as we have seen in the proof of Theorem 2.1. Conversely, suppose that $M \in \mathcal{M}'_A$ and $M_1 = M^G \otimes_{A^G} A$. The canonical map $\varphi : M_1 \rightarrow M$ is a morphism in $\mathcal{M}_A$. It is surjective by the definition of $\mathcal{M}'_A$. Now $\ker \varphi \in \mathcal{M}'_A$ by (1), hence $\ker \varphi$ is generated over A by G -invariant elements. But we have proved already that $M_1^G = M^G \otimes 1$. Since φ is injective on M_1^G , we get $\ker \varphi = 0$, i.e., φ is an isomorphism.

(3) follows from [1], Ch. I, §3, Prop. 12 and Ch. II, §5, Prop. 4. $\square$

Suppose that $M \in \mathcal{M}_A$ is finitely generated over A . Put

$$\operatorname{rk} M = \dim_K M \otimes_A K \quad \text{and} \quad M(x) = M/M\mathfrak{m}_x \quad \text{for } x \in X$$

where $\mathfrak{m}_x$ is the maximal ideal of A corresponding to x . Then $\dim M(x) \geq \operatorname{rk} M$ for all $x \in X$. By [1], Ch. 2, §3, Prop. 7 and §5, Corollary to Prop. 2, the set

$$X_M = \{x \in X \mid \dim M(x) = \operatorname{rk} M\}$$

is open in X and consists precisely of those x for which $M_{\mathfrak{m}_x}$ is a free $A_{\mathfrak{m}_x}$ -module. If $\mathfrak{m}_x$ is stable under G then $\mu^M(M\mathfrak{m}_x) = \mu^M(M) \cdot \mu^*(\mathfrak{m}_x) \subset M\mathfrak{m}_x \otimes R$ since $\mu^*(\mathfrak{m}_x) \subset \mathfrak{m}_x \otimes R$. In general, applying this observation to the action of G_x , we see that $M\mathfrak{m}_x$ is stable under G_x , and so G_x operates in $M(x)$. Put

$$s(M) = \min_{x \in X_{G\text{-reg}} \cap X_M} \dim M(x)^{G_x},$$

$$X_{M\text{-reg}} = \{x \in X_{G\text{-reg}} \mid \dim M(x) = \operatorname{rk} M \quad \text{and} \quad \dim M(x)^{G_x} = s(M)\}.$$

We call $X_{M\text{-reg}}$ the set of M -regular points in X .

THEOREM 3.3. (1) $X_{M\text{-reg}}$ is a G -invariant open subset of X .

(2) For all $x \in X_{M\text{-reg}}$ the canonical map $M \rightarrow M(x)$ induces a surjection $M^G \rightarrow M(x)^{G_x}$.

(3) If $X_{M\text{-reg}} = X$ then the map $M^G \otimes_{A^G} A \rightarrow M$ given by $m \otimes a \mapsto ma$ is a split monomorphism of A -modules and M^G is projective of rank $s(M)$ over A^G .

(4) $\dim_{K^G}(M \otimes_A K)^G = s(M)$.

Proof. If $x \in X$ and $g \in G(k)$ then g induces a linear isomorphism $M(x) \rightarrow M(xg)$ compatible with the actions of stabilizers. Hence $M(x)^{G_x} \cong M(xg)^{G_{xg}}$. It follows that X_M and $X_{M\text{-reg}}$ are stable under the action of $G(k)$. Then X_M is a G -invariant open subset, and so is $X_{G\text{-reg}} \cap X_M$ too. Localizing if necessary, we may assume that $X = X_{G\text{-reg}}$ and $M_{\mathfrak{m}_x}$ is free over $A_{\mathfrak{m}_x}$ for all x . Then M is a projective A -module.

We are going to apply Lemma 1.3 in which we take $F = M \otimes R$ with the A -module structure obtained again via $p_1^* : A \rightarrow A \otimes R$. Take $F'' = M \otimes 1$, which is clearly a direct summand of F . Put $F' = \mu^M(M) \cdot (A \otimes 1)$, and let $I'(x)$, $I''(x)$ be the images, respectively, of $F'/F'\mathfrak{m}_x$ and $F''/F''\mathfrak{m}_x$ in $F/F\mathfrak{m}_x \cong M(x) \otimes R$. We have $I''(x) \cong M(x) \otimes 1$. If μ_x^M denotes the composite

$$M \xrightarrow{\mu^M} M \otimes R \xrightarrow{\operatorname{can} \otimes \operatorname{id}_R} M(x) \otimes R,$$

then $I'(x) = \mu_x^M(M)$. Consider two G -module structures on $M \otimes R$: the first one is the tensor product of the given G -module structure on M and the left regular G -module structure on R ; the second one is the tensor product of the trivial G -module

structure on M and the right regular G -module structure on R . The map μ^M is G -equivariant with respect to the second structure and is a bijection of M onto the subspace $(M \otimes R)^G$ of G -invariant elements with respect to the first structure (see [9], Part I, 3.7, (5) and (6); however, we interchanged the left and right regular G -module structures). These two structures on $M \otimes R$ induce a G_x -module and a G -module structures on $M(x) \otimes R$. We get

$$\mu_x^M(M) \subset (M(x) \otimes R)^{G_x} = \text{ind}_{G_x}^G M(x).$$

Since μ_x^M is G -equivariant, $\mu_x^M(M)$ is a G -submodule of the induced module. Furthermore, $\mu_x^M(ma) = \mu_x^M(m) \cdot \mu_x^*(a)$ for $m \in M$, $a \in A$, whence $\mu_x^M(M)$ is stable under the action of $\mu_x^*(A) \subset R$. By Proposition 1.1 $\mu_x^*(A) = k[G_x \backslash G]$. The canonical G_x -equivariant map $\varphi : \text{ind}_{G_x}^G M(x) \rightarrow M(x)$ is the restriction of the map $\text{id} \otimes \varepsilon : M(x) \otimes R \rightarrow M(x)$. Hence the composite $\varphi \circ \mu_x^M$ coincides with the canonical projection $M \rightarrow M(x)$, it is therefore surjective. It follows that the inclusion $\iota : \mu_x^M(M) \hookrightarrow \text{ind}_{G_x}^G M(x)$ corresponds under the equivalence of the Imprimitivity Theorem to a surjective map of G_x -modules. Then ι must itself be surjective, i.e., $\mu_x^M(M) = \text{ind}_{G_x}^G M(x)$. In particular,

$$\dim I'(x) = \dim \text{ind}_{G_x}^G M(x) = (G : G_x) \cdot \dim M(x) = q(X) \text{rk}(M),$$

which does not depend on x . By Lemma 1.2 F' is a direct summand of F . Thus the hypotheses of Lemma 1.3 are fulfilled. By our previous description $I'(x) \cap I''(x) = (M(x) \otimes 1)^{G_x} \cong M(x)^{G_x}$. Hence $s = s(M)$ and $X_{M\text{-reg}}$ is the set of rational points of the open subset $U \subset \text{Spec } A$ defined in Lemma 1.3. Thus $X_{M\text{-reg}}$ is open in X . It is G -invariant by observation at the beginning of the proof. Therefore we may localize further and assume $X = X_{M\text{-reg}}$. By Lemma 3.1 $F' \cap F'' = M^G A \otimes 1$. Then (2) of Lemma 1.3 implies (2) of the theorem. Lemma 1.3 ensures that the A -module $M^G A$ is projective of rank $s(M)$ and is a direct summand of the A -module M . By Proposition 3.2 M^G is a projective A^G -module of rank $s(M)$ and $M^G \otimes_{A^G} A$ is mapped isomorphically onto $M^G A$. This proves (3). Assertion (4) is immediate since $(M \otimes_A K)^G \cong M^G \otimes_{A^G} K^G$. $\square$

Let $M, N \in \mathcal{M}_A$ and $P = \text{Hom}_A(N, M)$. For every finite dimensional commutative algebra B we have

$$P \otimes B \cong \text{Hom}_{A \otimes B}(N \otimes B, M \otimes B).$$

If $g \in G(B)$ and $\xi \in P \otimes B$ then we put $g_P(\xi) = g_M \circ \xi \circ g_N^{-1}$ where g_M and g_N are the operators on $M \otimes B$ and $N \otimes B$, respectively, corresponding to g . In this way we obtain a group action of $G(B)$ on $P \otimes B$ which is natural in B . If B is infinite dimensional then each point $g \in G(B)$ still belongs to $G(B')$ where $B' \subset B$ is a finite dimensional subalgebra. Indeed, we can take B' to be the image of the algebra homomorphism $R \rightarrow B$ corresponding to g . Extend the action of g in $P \otimes B'$ by B -linearity to the action in $P \otimes B$. If $g, h \in G(B)$ are two points then there exists a finite dimensional subalgebra B' such that $G(B')$ contains both of them. It follows that $(gh)_P = g_P h_P$. Thus P is equipped with a G -module structure, which is clearly compatible with the A -module structure, i.e., $P \in \mathcal{M}_A$.

Let V be a G -module. Then $V \otimes A$, considered with the natural A -module structure and the tensor product G -module structure, is an object of $\mathcal{M}_A$. Hence so is $\text{Hom}(V, M) \cong \text{Hom}_A(V \otimes A, M)$ too.

Suppose that $M, N \in \mathcal{M}_A$ are finitely generated over A and V a finite dimensional G -module. Put

$$s(N, M) = \min_{x \in X_{G\text{-reg}} \cap X_M \cap X_N} \dim \text{Hom}_{G_x}(N(x), M(x)),$$

$$s(V, M) = \min_{x \in X_{G\text{-reg}} \cap X_M} \dim \text{Hom}_{G_x}(V, M(x)).$$

COROLLARY 3.4. (1) $\dim_{K^G} \text{Hom}_{(K, G)}(N \otimes_A K, M \otimes_A K) = s(N, M)$.

(2) $\dim_{K^G} \text{Hom}_G(V, M \otimes_A K) = s(V, M)$.

(3) $\dim_{K^G} \text{soc}_G(M \otimes_A K) = \sum s(V, M) \dim V$, the sum over isomorphism classes of irreducible G -modules V .

Proof. (1) Let $P = \text{Hom}_A(N, M)$. Then $P \otimes_A K \cong \text{Hom}_K(N \otimes_A K, M \otimes_A K)$ and $P(x) \cong \text{Hom}(N(x), M(x))$ for all $x \in X_N$. Since $X_M \cap X_N \subset X_P$, we get $s(P) = s(N, M)$. Apply Theorem 3.3(4) to the (A, G) -module P , noting that

$$\text{Hom}_{(K, G)}(N \otimes_A K, M \otimes_A K) \cong (P \otimes_A K)^G.$$

Assertion (2) is a special case of (1) with $N = V \otimes A$. Next, the G -socle $\text{soc}_G(M \otimes_A K)$ is a direct sum of isotypic components I_V corresponding to irreducible G -modules V . Since $\text{End}_G V \cong k$, we have $I_V \cong \text{Hom}_G(V, M \otimes_A K) \otimes V$, and (3) follows from (2). $\square$

We continue to assume that $M \in \mathcal{M}_A$ is finitely generated over A .

LEMMA 3.5. If $x \in X_{G\text{-reg}} \cap X_M$ and $\mathfrak{n} = \mathfrak{m}_x \cap A^G$ then $M/M\mathfrak{n} \cong \text{ind}_{G_x}^G M(x)$ as G -modules. The restriction of the canonical map $M/M\mathfrak{n} \rightarrow M(x)$ yields a linear isomorphism $(M/M\mathfrak{n})^G \cong M(x)^{G_x}$.

Proof. Since $X_{G\text{-reg}} \cap X_M$ is a G -invariant open subset, there exists $f \in A^G \setminus \mathfrak{n}$ such that A_f is free of rank $q(X)$ over A_f^G and M_f is free over A_f . Then M_f is free of rank $q(X) \text{rk}(M)$ over A_f^G , and so $M/M\mathfrak{n} \cong M_f/M_f\mathfrak{n}$ has dimension $q(X) \text{rk}(M)$. In the proof of Theorem 3.3 we constructed a surjective G -module homomorphism $\mu_x^M : M \rightarrow \text{ind}_{G_x}^G M(x)$. Since $\mu^M(M\mathfrak{n}) = \mu^M(M) \cdot (\mathfrak{n} \otimes 1) \subset M\mathfrak{m}_x \otimes R$, we see that $M\mathfrak{n} \subset \ker \mu_x^M$. Comparing dimensions, we conclude that $M\mathfrak{n} = \ker \mu_x^M$. The final assertion is a special case of the Frobenius reciprocity. $\square$

Put $X_{M\text{-inj}} = \{x \in X_{G\text{-reg}} \cap X_M \mid M(x) \text{ is an injective } G_x\text{-module}\}$.

THEOREM 3.6. (1) $X_{M\text{-inj}}$ is open in X and consists precisely of those points $x \in X_{G\text{-reg}} \cap X_M$ for which there exists $f \in A^G$ such that $f(x) \neq 0$ and M_f is a projective (A_f, G) -module.

(2) $X_{M\text{-inj}}$ is nonempty if and only if $M \otimes_A K$ is a projective (K, G) -module.

(3) The induced G -modules $\text{ind}_{G_x}^G M(x)$ corresponding to points $x \in X_{M\text{-inj}}$ are all isomorphic to each other.

(4) For every $x \in X_{M\text{-inj}}$ there exist $f \in A^G$ and a G -submodule $V \subset M$ such that $f(x) \neq 0$, $V \cong \text{ind}_{G_x}^G M(x)$ and the linear map $V \otimes A_f^G \rightarrow M_f$ given by the rule $m \otimes a \mapsto ma$ is bijective. In particular, M_f is an injective G -module.

(5) If $X_{M\text{-inj}}$ is nonempty then $M \otimes_A K \cong V \otimes K^G$ as (K^G, G) -modules. In particular, $M \otimes_A K$ is an injective G -module.

Proof. Let $x \in X_{M\text{-inj}}$. The G -module $V_x = \text{ind}_{G_x}^G M(x)$ is injective by [9], Part I, 3.9, hence it is also projective. Then, in view of Lemma 3.5, there exists a G -submodule $V \subset M$ such that $M = V \oplus M\mathfrak{n}$ where $\mathfrak{n} = \mathfrak{m}_x \cap A^G$. Clearly $V \cong V_x$. Define a homomorphism of A^G -modules $\varphi : V \otimes A^G \rightarrow M$ by the formula $\varphi(m \otimes a) = ma$ for $m \in V$, $a \in A^G$. Since $x \in X_{G\text{-reg}} \cap X_M$, the algebra $A_{\mathfrak{n}}$ is free over $A_{\mathfrak{n}}^G$ and $M_{\mathfrak{n}}$ is free over $A_{\mathfrak{n}}$. Then $M_{\mathfrak{n}}$ is free over $A_{\mathfrak{n}}^G$, and it follows that φ induces an isomorphism $V \otimes A_{\mathfrak{n}}^G \rightarrow M_{\mathfrak{n}}$. There exists $f \in A^G \setminus \mathfrak{n}$ such that $\varphi_f : V \otimes A_f^G \rightarrow M_f$ is an isomorphism (see [1], Ch. 3, §5, Prop. 2). As a G -module, M_f is a direct sum of a family of copies of the G -module V , and so it is injective. This proves (4), and (5) is an immediate consequence.

If $\mathfrak{n}'$ is a maximal ideal of A^G such that $f \notin \mathfrak{n}'$ then $A_f^G = k \oplus A_f^G \mathfrak{n}'$, whence $M_f = V \oplus M_f \mathfrak{n}'$, and $M/M\mathfrak{n}' \cong M_f/M_f \mathfrak{n}' \cong V$ as G -modules. Again by Lemma 3.5 $\text{ind}_{G_y}^G M(y) \cong V$ for all $y \in X_{G\text{-reg}} \cap X_M$ such that $f(y) \neq 0$. If $x' \in X_{M\text{-inj}}$ is another point then, similarly, $\text{ind}_{G_y}^G M(y) \cong V_{x'}$ for all y in a nonempty open subset of X . Since X is irreducible, we conclude $V_x \cong V_{x'}$, whence (3).

Suppose that $N \in \mathcal{M}_A$ is free of finite rank over A and $\varphi : N \rightarrow M$ is an epimorphism in $\mathcal{M}_A$. Let $\varphi_x : N(x) \rightarrow M(x)$ be the epimorphism of G_x -modules obtained from φ by reduction modulo $\mathfrak{m}_x$. Since $M(x)$ is projective, there exists a G_x -module homomorphism $\psi_x : M(x) \rightarrow N(x)$ such that $\varphi_x \circ \psi_x = \text{id}_{M(x)}$. Put $P = \text{Hom}_A(M, N)$. Then $X_M \subset X_P$ and for $x \in X_M$ there is an isomorphism of G_x -modules $P(x) \cong \text{Hom}(M(x), N(x)) \cong M(x)^* \otimes N(x)$. Since $k[G_x]^*$ is a Frobenius algebra, the G_x -module $M(x)^*$ is injective. By [9], Part I, 3.10 $P(x)$ is also injective. As we know already, there exists a G -submodule $W \subset P$ such that $P = W \oplus P\mathfrak{n}$. By Lemma 3.5 the restriction of the canonical map $W \xrightarrow{\sim} P/P\mathfrak{n} \rightarrow P(x)$ yields a linear isomorphism $W^G \xrightarrow{\sim} P(x)^{G_x}$. Let $\psi \in W^G \subset P^G$ be the element corresponding to $\psi_x \in P(x)^{G_x}$. We may regard ψ as a morphism $M \rightarrow N$ in $\mathcal{M}_A$ whose reduction modulo $\mathfrak{m}_x$ is ψ_x . Then $\gamma = \varphi \circ \psi$ is an $\mathcal{M}_A$ -endomorphism of M whose reduction modulo $\mathfrak{m}_x$ is the identity transformation of $M(x)$. Let U be the set of those $y \in X_M$ for which the reduction of γ modulo $\mathfrak{m}_y$ is invertible. By [1], Ch. 2, §3, Corollary to Prop. 6 and §5, Prop. 2 U is open and consists precisely of those $y \in X_M$ for which $\gamma_{\mathfrak{m}_y} : M_{\mathfrak{m}_y} \rightarrow M_{\mathfrak{m}_y}$ is bijective. Since γ is $G(k)$ -equivariant, U is G -invariant. As $U = \pi^{-1}(\pi(U))$ and $\pi(U)$ is an open neighbourhood of $\pi(x)$ in X/G , there exists $f \in A^G$ such that $x \in X_f \subset U$ where $X_f = \{y \in X \mid f(y) \neq 0\}$. Then $\gamma_f : M_f \rightarrow M_f$ is bijective, and therefore $N_f = \ker \varphi_f \oplus \text{im } \psi_f$. In other words, $\varphi_f : N_f \rightarrow M_f$ is a split epimorphism in $\mathcal{M}_{A_f}$. Since M is finitely generated over A , it is an epimorphic image of a finitely generated free $A \# k[G]^*$ -module. We can take the latter to be our N . We see that M_f is a direct summand of a free $A_f \# k[G]^*$ -module for a suitable f .

Conversely, suppose that x is any point in $X_{G\text{-reg}} \cap X_M \cap X_f$ such that M_f is a projective $A_f \# k[G]^*$ -module. If $\mathfrak{n} = \mathfrak{m}_x \cap A^G$ then $M/M\mathfrak{n} \cong M_f/M_f \mathfrak{n}$ is a projective $(A/A\mathfrak{n}) \# k[G]^*$ -module. By Theorem 2.1(2) $A/A\mathfrak{n} \cong k[G_x \setminus G]$. The G_x -module $M(x)$ corresponds to $M/M\mathfrak{n}$ under the category equivalence of the Imprimitivity Theorem. It is therefore projective, hence injective. We get (1).

Suppose that $M \otimes_A K$ is projective in $\mathcal{M}_K$. We want to show that M_f is projective in $\mathcal{M}_{A_f}$ for a suitable $0 \neq f \in A^G$ and then apply (1). Since X_M is open and G -invariant, we may assume that $X_M = X$ passing at the very beginning to a suitable localization of A . Then M is projective as an A -module. Let $\varphi : N \rightarrow M$ be an epimorphism in $\mathcal{M}_A$ with N a free $A \# k[G]^*$ -module. It extends to an epimorphism $\varphi_K : N \otimes_A K \rightarrow M \otimes_A K$ in $\mathcal{M}_K$. By our assumptions the latter

admits a splitting $\psi : M \otimes_A K \rightarrow N \otimes_A K$ in $\mathcal{M}_K$. Since N is free over A , the localizations N_f are identified with their images in $N \otimes_A K$, and the same is valid for M . Since M is finitely generated over A , hence also over A^G , we have $\psi(M) \subset N_f$ for a suitable f . Then $\psi(M_f) = N_f$, which means that $\varphi_f : N_f \rightarrow M_f$ is a split epimorphism in $\mathcal{M}_{A_f}$. That completes the proof of (2). $\square$

4. Actions with linearly reductive stabilizers.

We weaken our assumptions for this section considerably. In the next proposition G is any affine group scheme over k , not necessarily finite, and A is any G -algebra, not necessarily commutative. What we prove is a special case of results due to Doi [4] obtained in the context of coactions of Hopf algebras.

PROPOSITION 4.1. *The following properties of a G -algebra A are equivalent:*

- (1) *All objects $M \in \mathcal{M}_A$ are injective G -modules.*
- (2) *There exist an injective G -module Q and a homomorphism of G -modules $\psi : Q \rightarrow A$ such that $1 \in \psi(Q^G)$.*
- (3) *There exists a G -module homomorphism $\varphi : k[G] \rightarrow A$ (where $k[G]$ is given the left regular G -module structure) such that $\varphi(1) = 1$.*
- (4) *There are linear maps $\Phi_M : M \rightarrow M^G$, defined for each $M \in \mathcal{M}_A$, which are natural in M and satisfy $\Phi_M(m) = m$ for all $m \in M^G$.*
- (5) *The functor $M \mapsto M^G$ is exact on $\mathcal{M}_A$.*
- (6) *A is a projective (A, G) -module.*

If A is commutative they are equivalent to another property:

- (7) *Every object $M \in \mathcal{M}_A$ which is finitely generated and projective as an A -module is projective in $\mathcal{M}_A$.*

Proof. (1) $\Rightarrow$ (2). By (1) A is an injective G -module. So we can take $Q = A$ and $\psi = \text{id}_A$.

(2) $\Rightarrow$ (3). Let $q \in Q^G$ be an element such that $\psi(q) = 1$. By injectivity of Q the G -module homomorphism $k \rightarrow Q$ sending 1 to q extends to a homomorphism $k[G] \rightarrow Q$. Composing the latter with ψ , we get φ .

(3) $\Rightarrow$ (4). Define Φ_M as the composite map

$$M \xrightarrow{\mu^M} M \otimes k[G] \xrightarrow{\text{id}_M \otimes \varphi} M \otimes A \longrightarrow M$$

where the last map is afforded by the A -module structure on M . Recall that $\mu^M(M) = (M \otimes k[G])^G$. Since the two final maps in the decomposition of Φ_M are G -equivariant, we get $\Phi_M(M) \subset M^G$. If $m \in M^G$ then $\mu^M(m) = m \otimes 1$, whence $\Phi_M(m) = m$. That the maps Φ_M are natural in M is clear.

(4) $\Rightarrow$ (5). The fixed point functor is clearly left exact. Suppose that $\xi : M \rightarrow N$ is an epimorphism in $\mathcal{M}_A$. Given $n \in N^G$, take $m \in M$ such that $\xi(m) = n$. Then $\Phi_M(m) \in M^G$ and $\xi(\Phi_M(m)) = \Phi_N(\xi(m)) = n$. Thus ξ induces a surjection $M^G \rightarrow N^G$.

(5) $\Rightarrow$ (1). We may view $\text{Hom}(V, M) \cong V^* \otimes M$ for each finite dimensional G -module V as an (A, G) -module taking the tensor product of G -module structures and letting A operate on the second tensorand. If $W \subset V$ is a G -submodule then we have an epimorphism $\text{Hom}(V, M) \rightarrow \text{Hom}(W, M)$ in $\mathcal{M}_A$. Applying the fixed point functor, we deduce the surjectivity of the canonical map $\text{Hom}_G(V, M) \rightarrow$

$\text{Hom}_G(W, M)$. Since all G -modules are locally finite dimensional, this gives the injectivity of M .

(5) $\Leftrightarrow$ (6). Every morphism $A \rightarrow M$ in $\mathcal{M}_A$ is given by the rule $a \mapsto ma$ where $m \in M^G$. Hence $\text{Hom}_{(A,G)}(A, M) \cong M^G$. Note that the projectivity of A in $\mathcal{M}_A$ means that the functor $M \mapsto \text{Hom}_{(A,G)}(A, M)$ is exact.

(5) $\Rightarrow$ (7). If M, N are A -modules, $P = \text{Hom}_A(M, N)$ and B a commutative algebra then the canonical map $P \otimes B \rightarrow \text{Hom}_{A \otimes B}(M \otimes B, N \otimes B)$ is bijective when M is free of finite rank, hence also when M is finitely presented. If $M, N \in \mathcal{M}_A$ and M is finitely presented as an A -module then $G(B)$ operates in $P \otimes B$, naturally in B . This gives P a G -module structure. Assuming A to be commutative, we have $P \in \mathcal{M}_A$. If, moreover, M is projective as an A -module then every epimorphism $N \rightarrow N'$ in $\mathcal{M}_A$ induces an epimorphism $\text{Hom}_A(M, N) \rightarrow \text{Hom}_A(M, N')$. Applying the fixed point functor, we deduce the surjectivity of the map $\text{Hom}_{(A,G)}(M, N) \rightarrow \text{Hom}_{(A,G)}(M, N')$.

(7) $\Rightarrow$ (6) is obvious. $\square$

Let X be an arbitrary scheme over k , and G a finite group scheme operating on X from the right. We still need the assumption that X can be covered by G -invariant affine open subschemes. We say that the stabilizer G_x of a point $x \in X(k)$ is *linearly reductive* if all G_x -modules are completely reducible. This is equivalent to the semisimplicity of the Hopf algebra $k[G_x]^*$. By [3], Ch. IV, §3, 3.6 G_x is linearly reductive if and only if its identity component G_x^0 is diagonalizable and the index $(G_x : G_x^0)$ is prime to $p = \text{char } k$ when $p > 0$. Put

$$X_{\text{lin.red.}} = \{x \in X(k) \mid G_x \text{ is linearly reductive}\}.$$

THEOREM 4.2. *The set $X_{\text{lin.red.}}$ consists precisely of those $x \in X(k)$ which are contained in a G -invariant affine open subscheme $U \subset X$ such that $k[U]$ is an injective G -module. In particular, $X_{\text{lin.red.}}$ is the set of rational points of an open G -invariant subscheme of X . If, moreover, X is an algebraic variety, then:*

(1) *The condition $X_{\text{lin.red.}} \neq \emptyset$ is equivalent to each of the two below:*

(a) *$k(X)$ is an injective G -module.*

(b) *The smash product algebra $k(X) \# k[G]^*$ is semisimple.*

(2) *For every $x \in X_{G\text{-reg}} \cap X_{\text{lin.red.}}$ there exist a G -invariant affine open neighbourhood U of x and a G -submodule $V \subset k[U]$ such that $V \cong \text{ind}_{G_x}^G k$ and the map $V \otimes k[U]^G \rightarrow k[U]$ given by the multiplication in $k[U]$ is bijective.*

(3) *If $X_{\text{lin.red.}}$ is nonempty then there exists a G -submodule $V \subset k[X]$ such that $V \cong \text{ind}_{G_x}^G k$ for all $x \in X_{G\text{-reg}} \cap X_{\text{lin.red.}}$ and the map $V \otimes k(X)^G \rightarrow k(X)$ given by the multiplication in $k(X)$ is bijective.*

Proof. We may assume that X is affine. Put $A = k[X]$. Given $x \in X(k)$, the orbit morphism $\mu_x : G \rightarrow X$ determines a G -equivariant homomorphism of algebras $\mu_x^* : A \rightarrow k[G]$ whose image is $k[G_x \backslash G]$ by Proposition 1.1. If $x \in X_{\text{lin.red.}}$ then all G_x -modules are injective. Hence $k[G_x \backslash G] = \text{ind}_{G_x}^G k$ is an injective G -module by [9], Part I, 3.9. As it is also projective, there is a G -submodule $V \subset A$ mapped isomorphically onto $k[G_x \backslash G]$ under μ_x^* . Take $f \in V$ such that $\mu_x^*(f) = 1$. Then $f \in A^G$, and the map $V \rightarrow A_f$, $v \mapsto vf^{-1}$, is a G -module homomorphism under which $f \mapsto 1$. Since V is an injective G -module, so is A_f too by implication (2) $\Rightarrow$ (1) of Proposition 4.1. Furthermore, $f(x) = 1$ since $\ker \mu_x^* \subset \mathfrak{m}_x$. Thus $\text{Spec } A_f$ is the required open neighbourhood of x .

Conversely, suppose that $x \in U(k)$ where $U \subset X$ is a G -invariant affine open subscheme such that $k[U]$ is an injective G -module. As μ_x factors through U , it induces a G -equivariant algebra homomorphism $k[U] \rightarrow k[G_x \backslash G]$. Hence we may view $k[G_x \backslash G]$ as a $(k[U], G)$ -module. By Proposition 4.1 $k[G_x \backslash G]$ is an injective G -module. It is then a direct summand of $k[G]$. By [9], Part I, 4.12 $k[G_x \backslash G]$ is also an injective G_x -module. Now k is a direct summand of $k[G_x \backslash G]$ as a G_x -module. Hence k is an injective G_x -module. Then all G_x -modules are injective, which implies that all G_x -modules are completely reducible.

Suppose that X is an algebraic variety and $K = k(X)$. Apply Theorem 3.6 to the (A, G) -module A . Noting that $X_{\text{lin.red.}}$ is precisely the set of points x for which $A/\mathfrak{m}_x \cong k$ is an injective G_x -module, we get assertions (2) and (3) of Theorem 4.2. Furthermore, $X_{\text{lin.red.}}$ is nonempty if and only if K is a projective (K, G) -module. By Proposition 4.1 this is equivalent to K being an injective G -module. This is equivalent also to the condition that every $M \in \mathcal{M}_K$ of finite dimension over K is projective in $\mathcal{M}_K$. This means, in particular, that all ideals of the algebra $K \# k[G]^*$ are projective, which is equivalent to condition (b). $\square$

5. Invariants of restricted Lie algebras.

Suppose that $\text{char } k = p > 0$. Let X be an affine algebraic variety, and $\mathfrak{g}$ a p -Lie algebra over k . Put $A = k[X]$ and $K = k(X)$. Define an *action* of $\mathfrak{g}$ on X to be a homomorphism of p -Lie algebras $\rho : \mathfrak{g} \rightarrow \text{Der } A$ into the derivation algebra of A . Define $\mathfrak{g}_x \subset \mathfrak{g}$ to be the stabilizer of the maximal ideal $\mathfrak{m}_x$ of A corresponding to a point $x \in X$. Since $\rho(\mathfrak{g})(\mathfrak{m}_x^2) \subset \mathfrak{m}_x$, we have a linear map $\mathfrak{g} \rightarrow T_x X = (\mathfrak{m}_x/\mathfrak{m}_x^2)^*$ whose kernel is precisely $\mathfrak{g}_x$. Hence $\text{codim}_{\mathfrak{g}} \mathfrak{g}_x \leq \dim T_x X$. Since the dimensions of tangent spaces are bounded, it is meaningful to define

$$c_{\mathfrak{g}}(X) = \max_{x \in X} \text{codim}_{\mathfrak{g}} \mathfrak{g}_x,$$

$$X_{\mathfrak{g}\text{-reg}} = \{x \in X \mid \text{codim}_{\mathfrak{g}} \mathfrak{g}_x = c_{\mathfrak{g}}(X)\}.$$

If $\dim \mathfrak{g} < \infty$, there is a finite group scheme of height one $G = \mathfrak{G}(\mathfrak{g})$ associated with $\mathfrak{g}$ (see [3], Ch. II, §7, 3.9). One has $k[G] \cong u(\mathfrak{g})^*$ where $u(\mathfrak{g})$ is the restricted universal enveloping algebra of $\mathfrak{g}$. The action of $\mathfrak{g}$ on X corresponds to a group action of G according to [3], Ch. II, §7, 3.10. Furthermore, $G_x \cong \mathfrak{G}(\mathfrak{g}_x)$, so that $k[G_x \backslash G] \cong \text{Hom}_{u(\mathfrak{g}_x)}(u(\mathfrak{g}), k)$ and $(G : G_x) = p^{\text{codim}_{\mathfrak{g}} \mathfrak{g}_x}$ for all $x \in X$. It follows then that $X_{\mathfrak{g}\text{-reg}} = X_{G\text{-reg}}$. However, we want to extend Theorem 2.1 to the case of infinite dimensional $\mathfrak{g}$.

The Lie algebra $\text{Der } A$ has a natural A -module structure. Given $f \in A$ and $D, D' \in \text{Der } A$, we have

$$[fD, D'] = f[D, D'] - D'(f)D,$$

$$(fD)^p = f^p D^p + (fD)^{p-1}(f)D.$$

The first formula is easily checked straightforwardly. The second one is proved by Hochschild [8], Lemma 1. It follows that the A -submodule $L = A \cdot \rho(\mathfrak{g})$ is also a p -Lie subalgebra of $\text{Der } A$. Define a linear map

$$d : A \rightarrow L_A^* = \text{Hom}_A(L, A)$$

by the rule $(df)(D) = D(f)$ for $f \in A$, $D \in L$.

LEMMA 5.1. (1) $c_{\mathfrak{g}}(X) = c_L(X)$ and $X_{\mathfrak{g}\text{-reg}} = X_{L\text{-reg}}$.

(2) $X_{L\text{-reg}}$ is open in X and consists precisely of those $x \in X$ for which $L_{\mathfrak{m}_x}$ is a free $A_{\mathfrak{m}_x}$ -module and $L_A^* = dA + \mathfrak{m}_x L_A^*$.

(3) If $X_{L\text{-reg}} = X$ then L is projective of rank $c_L(X)$ over A and $L_A^* = A \cdot dA$.

Proof. If L_x is the stabilizer of $\mathfrak{m}_x$ in L then $\mathfrak{m}_x L \subset L_x$, whence $L = \rho(\mathfrak{g}) + L_x$. It follows that $L/L_x \cong \mathfrak{g}/\mathfrak{g}_x$, and (1) is immediate.

Take a finite system of generators $a_1, \dots, a_n$ of the algebra A and define a homomorphism of A -modules $\varphi : L \rightarrow F$, where $F = A^n$, by the rule $\varphi(D) = (Da_1, \dots, Da_n)$ for $D \in L$. Since each derivation of A is determined by its values on generators, we have $\ker \varphi = 0$. Hence $L \cong F' = \varphi(L) \subset F$. Denote by $I(x)$ the image of the map $L/\mathfrak{m}_x L \cong F'/\mathfrak{m}_x F' \rightarrow F/\mathfrak{m}_x F$ induced by φ . If $D \in L$ then $D \in L_x$ if and only if $D(A) \subset \mathfrak{m}_x$ (as $A = k + \mathfrak{m}_x$), if and only if $Da_i \in \mathfrak{m}_x$ for all i . It follows that $I(x) \cong L/L_x$. Applying Lemma 1.2, we see that $X_{L\text{-reg}}$ is open and coincides with the set of points $x \in X$ for which $F'_{\mathfrak{m}_x}$ is a direct summand of the $A_{\mathfrak{m}_x}$ -module $F_{\mathfrak{m}_x}$, i.e., $\varphi_{\mathfrak{m}_x} : L_{\mathfrak{m}_x} \rightarrow F_{\mathfrak{m}_x}$ is a split monomorphism of $A_{\mathfrak{m}_x}$ -modules. Assuming that the $A_{\mathfrak{m}_x}$ -module $L_{\mathfrak{m}_x}$ is free, $\varphi_{\mathfrak{m}_x}$ splits if and only if the localization at $\mathfrak{m}_x$ of the dual homomorphism $\varphi^* : \text{Hom}_A(F, A) \rightarrow \text{Hom}_A(L, A)$ is surjective. By Nakayama's lemma this is equivalent to the equality $L_A^* = N + \mathfrak{m}_x L_A^*$ where N is the image of φ^* . Clearly, N is the A -submodule of L_A^* generated by $da_1, \dots, da_n$. Since $d(ab) = a \cdot db + b \cdot da$ for $a, b \in A$, we have $N = A \cdot dA$. Then $N + \mathfrak{m}_x L_A^* = dA + \mathfrak{m}_x L_A^*$ since $A = k + \mathfrak{m}_x$. We get (2).

If $x \in X_{L\text{-reg}}$ then the map $L/\mathfrak{m}_x L \rightarrow F/\mathfrak{m}_x F$ is injective. Hence $L_x = \mathfrak{m}_x L$ and $\dim L/\mathfrak{m}_x L = c_L(X)$. Now (3) follows from (2) by globalization. $\square$

THEOREM 5.2. The subset $X_{\mathfrak{g}\text{-reg}}$ is open in X . Furthermore:

(1) If $X_{\mathfrak{g}\text{-reg}} = X$ then A is a projective $A^{\mathfrak{g}}$ -module of rank $p^{c_{\mathfrak{g}}(X)}$.

(2) If $x \in X_{\mathfrak{g}\text{-reg}}$ and $\mathfrak{n} = \mathfrak{m}_x \cap A^{\mathfrak{g}}$ then $\mathfrak{n}A$ is a maximal $\mathfrak{g}$ -invariant ideal of A and the algebra $A/\mathfrak{n}A$ is $\mathfrak{g}$ -equivariantly isomorphic with $\text{Hom}_{u(\mathfrak{g}_x)}(u(\mathfrak{g}), k)$.

(3) If $X_{\mathfrak{g}\text{-reg}} = X$ then the assignment $I \mapsto I^{\mathfrak{g}}$ establishes a bijection between the $\mathfrak{g}$ -invariant ideals of A and the ideals of $A^{\mathfrak{g}}$. The canonical maps $A^{\mathfrak{g}} \rightarrow (A/I)^{\mathfrak{g}}$ are surjective.

(4) If $X_{\mathfrak{g}\text{-reg}} = X$ then $(B \otimes_{A^{\mathfrak{g}}} A)^{\mathfrak{g}} \cong B$ for every $A^{\mathfrak{g}}$ -algebra B on which $\mathfrak{g}$ operates trivially.

(5) $[k(X) : k(X)^{\mathfrak{g}}] = p^{c_{\mathfrak{g}}(X)}$.

Proof. As is immediate from the definition of L , the L -invariants coincide with the $\mathfrak{g}$ -invariants, and an ideal of A is stable under L if and only if it is stable under $\mathfrak{g}$. Since $\mathfrak{g}_x = \rho^{-1}(L_x)$, the algebra map $\text{Hom}_{u(L_x)}(u(L), k) \rightarrow \text{Hom}_{u(\mathfrak{g}_x)}(u(\mathfrak{g}), k)$ induced by ρ is an isomorphism. It follows that all assertions of the theorem for the p -Lie algebra $\mathfrak{g}$ are equivalent to corresponding assertions for the p -Lie algebra L . Put $c = c_{\mathfrak{g}}(X) = c_L(X)$.

Given $x \in X_{L\text{-reg}}$, we have $\dim L_A^*/\mathfrak{m}_x L_A^* = c$. Take $a_1, \dots, a_c \in A$ such that $da_1, \dots, da_c$ are a basis for a complement of $\mathfrak{m}_x L_A^*$ in L_A^* . Since the $A_{\mathfrak{m}_x}$ -module $L_{\mathfrak{m}_x}$ and its dual are free, passing to a suitable affine open neighbourhood of x , we may assume that L is a free A -module and $da_1, \dots, da_c$ are a basis for L_A^* over A . Let $D_1, \dots, D_c$ be the dual basis for L over A . This means that $D_i a_l = da_l(D_i) = \delta_{il}$ for all i, l . As L is a Lie subalgebra, we have $[D_i, D_j] = \sum_{l=1}^c g_{ijl} D_l$ for certain $g_{ijl} \in A$. Applying the derivations on both sides of the equality to a_l , we deduce $g_{ijl} = 0$. Since L is closed under p -th powers, we have $D_i^p = \sum_{l=1}^c h_{il} D_l$ for certain $h_{il} \in A$.

We deduce similarly that $h_{il} = 0$. Thus the linear span $\mathfrak{a} = \langle D_1, \dots, D_c \rangle \subset L$ is an abelian Lie subalgebra with zero p -map. Since $L = A\mathfrak{a}$, the assertions of the theorem for L are equivalent to those for $\mathfrak{a}$. Since $\dim \mathfrak{a} < \infty$ they are equivalent also to the assertions of Theorem 2.1 for the corresponding action of the finite group scheme $\mathfrak{G}(\mathfrak{a})$ (in fact this action is free). $\square$

COROLLARY 5.3. *Suppose that $\mathfrak{h} \subset \mathfrak{g}$ is a p -Lie subalgebra such that $\mathfrak{g} = \mathfrak{h} + \mathfrak{g}_x$ for at least one $x \in X_{\mathfrak{g}\text{-reg}}$. Then $A^{\mathfrak{g}} = A^{\mathfrak{h}}$.*

Proof. Obviously $\mathfrak{h}_x = \mathfrak{g}_x \cap \mathfrak{h}$. By the hypotheses $\mathfrak{h}/\mathfrak{h}_x \cong \mathfrak{g}/\mathfrak{g}_x$ for some point $x \in X_{\mathfrak{g}\text{-reg}}$. Then $c_{\mathfrak{h}}(X) \geq \text{codim}_{\mathfrak{h}} \mathfrak{h}_x = c_{\mathfrak{g}}(X)$. It follows that $[K : K^{\mathfrak{h}}] \geq [K : K^{\mathfrak{g}}]$. On the other hand, $K^{\mathfrak{g}} \subset K^{\mathfrak{h}}$, whence $K^{\mathfrak{g}} = K^{\mathfrak{h}}$. We conclude that $A^{\mathfrak{h}} = A \cap K^{\mathfrak{h}} = A \cap K^{\mathfrak{g}} = A^{\mathfrak{g}}$. $\square$

For every $r \geq 1$ put $A^{(p^r)} = \{f^{p^r} \mid f \in A\}$. The notations $K^{(p^r)}$, $\mathfrak{m}_x^{(p^r)}$ will have a similar meaning. For $f \in A$ let $d_x f : T_x X \rightarrow k$ denote the differential of f at x .

THEOREM 5.4. *Suppose that X is a smooth affine variety and $f_1, \dots, f_n \in A^{\mathfrak{g}}$ where $n = \dim X - c_{\mathfrak{g}}(X)$. Denote by U the open set of those $x \in X$ for which $d_x f_1, \dots, d_x f_n$ are linearly independent. If $U \neq \emptyset$ then $K^{\mathfrak{g}} = K^{(p)}(f_1, \dots, f_n)$. If, moreover, $\text{codim}_X X \setminus U \geq 2$ then:*

- (1) $A^{\mathfrak{g}} = A^{(p)}[f_1, \dots, f_n]$ and $A^{\mathfrak{g}}$ is free of rank p^n over $A^{(p)}$.
- (2) $A^{\mathfrak{g}}$ is a locally complete intersection.

(3) *If $\pi : X \rightarrow X/\mathfrak{g} = \text{Spec } A^{\mathfrak{g}}$ is the canonical morphism then $\pi(U)$ is the set of all smooth rational points of $X/\mathfrak{g}$.*

Proof. Put $B = A^{(p)}[f_1, \dots, f_n]$, $Y = \text{Spec } B$, $X^{(p)} = \text{Spec } A^{(p)}$. The scheme $X^{(p)}$ is obtained from X by base change $\mathfrak{f} : k \rightarrow k$ where $\mathfrak{f}$ is the Frobenius automorphism of k . Since smoothness is preserved under base change, $X^{(p)}$ is smooth. Denote by $\psi : Y \rightarrow X^{(p)}$ and $\varphi : X \rightarrow Y$ the morphisms corresponding to the inclusions $A^{(p)} \subset B \subset A$. Both φ and ψ are homeomorphisms. In particular, X , Y and $X^{(p)}$ have the same dimension. Put $d = \dim X$.

For each commutative algebra R denote by Ω_R the R -module of Kähler differentials of R over k . By [11], (27.B), $\dim_K \Omega_K = \deg \text{tr } K/k = d$ since K is separably generated over k . Furthermore, if $u_1, \dots, u_d \in K$ are any elements such that $du_1, \dots, du_d$ are a basis for Ω_K over K then the elements $u_1^{m_1} \dots u_d^{m_d}$ with $0 \leq m_i < p$ constitute a basis for K over $K^{(p)}$. In particular, $[K : K^{(p)}] = p^d$. We have $\Omega_K \cong \Omega_A \otimes_A K$ where Ω_A is a projective A -module since X is smooth (see [11], (29.B), Lemma 1). Assume that $U \neq \emptyset$. If $x \in U$ then $df_1, \dots, df_n$ are linearly independent modulo $\mathfrak{m}_x \Omega_A$, hence constitute a basis for a direct summand of the free $A_{\mathfrak{m}_x}$ -module $(\Omega_A)_{\mathfrak{m}_x}$. In particular, $df_1, \dots, df_n$ are linearly independent over A , hence also over K . It follows that $[L : K^{(p)}] = p^n$ where $L = K^{(p)}(f_1, \dots, f_n)$ is the field of fractions of B . Since $L \subset K^{\mathfrak{g}}$ and $[K : L] = p^{d-n} = p^{c_{\mathfrak{g}}(X)} = [K : K^{\mathfrak{g}}]$, we deduce $L = K^{\mathfrak{g}}$.

By the above B is free over its subalgebra $A^{(p)}$ with basis elements $f_1^{m_1} \dots f_n^{m_n}$ where $0 \leq m_i < p$. Then $B \cong A^{(p)}[t_1, \dots, t_n]/I$ where $t_1, \dots, t_n$ are indeterminates and I is the ideal of the polynomial algebra generated by n elements $t_i^p - f_i^p$. This means that Y is isomorphic with the scheme-theoretic fibre $F = \tau^{-1}(0)$ of the morphism $\tau : X^{(p)} \times \mathbb{A}^n \rightarrow \mathbb{A}^n$ where $\mathbb{A}^n$ is the affine space of dimension n and the components of τ are the functions $f_i^p - t_i^p$, $i = 1, \dots, n$. Note that F is a complete intersection in the smooth variety $X^{(p)} \times \mathbb{A}^n$ since F has codimension n . It follows that B is a locally complete intersection ring. In particular, B is Cohen-Macaulay.

The tangent space $T_z F$ at a point $z \in F$ coincides with the kernel of the linear map $d_z \tau : T_z(X^{(p)} \times \mathbb{A}^n) \rightarrow T_0 \mathbb{A}^n$ induced by τ in tangent spaces. Let $z = (x^{(p)}, a)$ where $x^{(p)} = (\psi \circ \varphi)(x) \in X^{(p)}$ for some $x \in X$ and $a \in \mathbb{A}^n$. It is easy to differentiate τ : for $(u, v) \in T_{x^{(p)}} X^{(p)} \oplus T_a \mathbb{A}^n \cong T_z(X^{(p)} \times \mathbb{A}^n)$ the vector $(d_z \tau)(u, v) \in T_0 \mathbb{A}^n \cong k^n$ has components $(d_{x^{(p)}} f_i^p)(u)$, $i = 1, \dots, n$. Note that $T_{x^{(p)}} X^{(p)} \cong T_x X \otimes_{\mathbb{F}} k$ and the maps $d_{x^{(p)}} f_i^p : T_x X \otimes_{\mathbb{F}} k \rightarrow k$ are just $d_x f_i \otimes \text{id}_k$ in this realization. The variety F is smooth at z if and only if $\dim T_z F = d$, if and only if $d_z \tau$ is surjective, if and only if $d_{x^{(p)}} f_1^p, \dots, d_{x^{(p)}} f_n^p$ are linearly independent, if and only if $d_x f_1, \dots, d_x f_n$ are linearly independent. In other words, the smoothness of Y at $\varphi(x)$ is equivalent to the inclusion $x \in U$. The codimension of the closed subset $Y \setminus \varphi(U)$ in Y is equal to that of $X \setminus U$ in X . Suppose that it is at least 2. Then Y is smooth in codimension 1. By Serre's normality criterion B is integrally closed (see [11], (17.I)). Then $A^{\mathfrak{g}} = B$ since both algebras have the same field of fractions. $\square$

Suppose that $\mathfrak{g} = \text{Lie } \mathcal{G}$ where $\mathcal{G}$ is a reduced algebraic group operating on X from the right. Then there is the induced action of $\mathfrak{g}$ on X . For $x \in X$ denote by $\mathcal{G}_x$ the scheme-theoretic stabilizer of x in $\mathcal{G}$. Let $X_{\text{reg}} \subset X$ be the open subset consisting of points x for which the orbit $x\mathcal{G}$ has a maximal possible dimension.

THEOREM 5.5. *Suppose that X is a smooth affine variety and $f_1, \dots, f_n \in A^{\mathcal{G}}$ where $n = \dim X - c_{\mathfrak{g}}(X)$. Suppose also that the open set U introduced in theorem 5.4 is nonempty. Then $X_{\mathfrak{g}\text{-reg}} = \{x \in X_{\text{reg}} \mid \mathcal{G}_x \text{ is reduced}\}$. If G denotes the r -th Frobenius kernel of $\mathcal{G}$ for some $r \geq 1$ then $K^G = K^{(p^r)}(f_1, \dots, f_n)$. If, moreover, $\text{codim}_X X \setminus U \geq 2$ then:*

- (1) $A^G = A^{(p^r)}[f_1, \dots, f_n]$ and A^G is free of rank p^{rn} over $A^{(p^r)}$.
- (2) A^G is a locally complete intersection.
- (3) $\pi(U)$ is the set of all smooth rational points of X/G where $\pi : X \rightarrow X/G$ is the canonical morphism.

Proof. Consider the morphism $\varphi : X \rightarrow \mathbb{A}^n$ with components $f_1, \dots, f_n$. Let $x \in U$. The differential $d_x \varphi : T_x X \rightarrow k^n$ is then surjective. By [3], Ch. I, §4, 4.15 φ is smooth, hence also flat at x . Then $\dim_x F_x = \dim X - \dim \mathbb{A}^n = d - n$ by [3], Ch. I, §3, 6.3, where $F_x = \varphi^{-1}(\varphi(x))$ and $d = \dim X$. Since F_x is a $\mathcal{G}$ -invariant closed subscheme of X , we have $x\mathcal{G} \subset F_x$, and so $\dim x\mathcal{G} \leq d - n = c_{\mathfrak{g}}(X)$.

Since U is open in X , we get $\dim x\mathcal{G} \leq c_{\mathfrak{g}}(X)$, hence also $\dim \mathcal{G}_x \geq \dim \mathcal{G} - c_{\mathfrak{g}}(X)$ for all $x \in X$. As $\dim \mathcal{G} = \dim \mathfrak{g}$, we can rewrite the last inequality in the form

$$c_{\mathfrak{g}}(X) - \text{codim}_{\mathfrak{g}} \mathfrak{g}_x \geq \dim \mathfrak{g}_x - \dim \mathcal{G}_x. \quad (*)$$

The subset X_{reg} consists of those x for which $\dim x\mathcal{G} = c_{\mathfrak{g}}(X)$, which is equivalent to an equality in $(*)$. By [3], Ch. III, §2, 2.6 $\text{Lie } \mathcal{G}_x = \mathfrak{g}_x$. Furthermore, $\dim \mathfrak{g}_x \geq \dim \mathcal{G}_x$ and the equality holds here precisely when $\mathcal{G}_x$ is smooth (which is equivalent to $\mathcal{G}_x$ being reduced for an algebraically closed field) by [3], Ch. II, §5, 2.1. As is easy to see, we have equalities everywhere above if and only if the left hand side of $(*)$ is zero, i.e., $x \in X_{\mathfrak{g}\text{-reg}}$. This proves the first assertion of the theorem.

We have $|G| = p^{r \dim \mathfrak{g}}$ (see [9], Part I, 9.6, (2)). Next, $G_x = G \cap \mathcal{G}_x$ for all $x \in X$. By [9], Part I, 9.4, (2) G_x coincides with the r -th Frobenius kernel of $\mathcal{G}_x$. If $x \in X_{\mathfrak{g}\text{-reg}}$ then $\mathcal{G}_x$ is reduced, whence $|G_x| = p^{r \dim \mathfrak{g}_x}$, and $(G : G_x) = |G|/|G_x| = p^{r \text{codim}_{\mathfrak{g}} \mathfrak{g}_x}$. Since both $X_{G\text{-reg}}$ and $X_{\mathfrak{g}\text{-reg}}$ are open and nonempty, they have a common point, which shows that $q(X) = p^{rc_{\mathfrak{g}}(X)}$ in the notations of Theorem 2.1. We deduce that $[K : K^G] = p^{rc_{\mathfrak{g}}(X)}$.

Let $L_i = K^{(p^i)}(f_1, \dots, f_n)$ for each $i \geq 1$. If $i > 1$ then $K^{(p^i)} \subset L_{i-1}^{(p)} \subset L_i$. As we have seen in Theorem 5.4 the elements $f_1^{m_1} \dots f_n^{m_n}$ with $0 \leq m_i < p$ are linearly independent over $K^{(p)}$. It follows that $[L_i : L_{i-1}^{(p)}] = p^n$. Hence

$$[L_i : K^{(p^i)}] = p^n [L_{i-1}^{(p)} : K^{(p^i)}] = p^n [L_{i-1} : K^{(p^{i-1})}].$$

We have also $[K : K^{(p^i)}] = [K : K^{(p)}] \cdot [K^{(p)} : K^{(p^i)}] = p^d [K : K^{(p^{i-1})}]$. Proceeding by induction on i we deduce that $[L_i : K^{(p^i)}] = p^{ni}$ and $[K : K^{(p^i)}] = p^{di}$. Taking $i = r$, we get $[K : L_r] = p^{(d-n)r} = [K : K^G]$. Since $L_r \subset K^G$, it follows $L_r = K^G$. The remainder of the theorem is proved in the same way as Theorem 5.4 with obvious changes. $\square$

Two classical cases of Theorem 5.5 are those when $\mathcal{G}$ is a semisimple algebraic group operating either on $\mathfrak{g}$ via the adjoint representation or on itself by conjugations. Let n denote the rank of $\mathcal{G}$. Under assumption that p does not divide the order of the Weyl group of $\mathcal{G}$ it was shown by Veldkamp [21] that $k[\mathfrak{g}]^{\mathcal{G}}$ is generated by n algebraically independent polynomials $J_1, \dots, J_n$ and $\mathfrak{g}_{\text{reg}}$ consists precisely of those $x \in \mathfrak{g}$ for which $d_x J_1, \dots, d_x J_n$ are linearly independent. The complement of $\mathfrak{g}_{\text{reg}}$ has pure codimension 3 in $\mathfrak{g}$. The stabilizer $\mathfrak{g}_x$ is just the centralizer of x in $\mathfrak{g}$. As $\dim \mathfrak{g}_x = n$ for all $x \in \mathfrak{g}_{\text{reg}}$, we have $c_{\mathfrak{g}}(\mathfrak{g}) = \dim \mathfrak{g} - n$.

Suppose that $\mathcal{G}$ is simply connected. Then the algebra of regular functions on $\mathcal{G}$, constant on the conjugacy classes, is generated by the characters $\chi_1, \dots, \chi_n$ of fundamental irreducible representations of $\mathcal{G}$. As shown by Steinberg [18], $\mathcal{G}_{\text{reg}}$ consists precisely of those $x \in \mathcal{G}$ for which $d_x \chi_1, \dots, d_x \chi_n$ are linearly independent. The complement of $\mathcal{G}_{\text{reg}}$ in $\mathcal{G}$ again has codimension 3. If $x \in \mathcal{G}$ then $\mathfrak{g}_x = \ker d_e \mu_x$ where $\mu_x : \mathcal{G} \rightarrow \mathcal{G}$ is the morphism defined by the rule $y \mapsto y^{-1}xy$. It follows $\mathfrak{g}_x = \{v \in \mathfrak{g} \mid (\text{Ad } x)v = v\}$. If $\mathcal{T} \subset \mathcal{G}$ is a maximal torus then there exists $t \in \mathcal{T}$ such that $\alpha(t) \neq 1$ for all roots α . Then $\mathfrak{g}_t = \text{Lie } \mathcal{T}$, and so $c_{\mathfrak{g}}(\mathcal{G}) = \dim \mathcal{G} - n$. No restrictions on p are needed in this case. The invariants of $\mathfrak{g}$ and Frobenius kernels were described by Friedlander and Parshall [6], and Donkin [5].

We give yet another example showing that Theorem 5.4 has a wider range of applications. Let $\mathfrak{g} = W_n$ be the Jacobson-Witt algebra. Recall that $\mathfrak{g} = \text{Der } B_n$ where $B_n = k[x_1, \dots, x_n]$, $x_i^p = 0$, is the truncated polynomial algebra. If $\mathcal{G} = \text{Aut } B_n$ then $\text{Lie } \mathcal{G}$ is the subalgebra of codimension n in $\mathfrak{g}$ consisting of derivations that leave stable the maximal ideal $\mathfrak{n}$ of B_n . For $D \in \mathfrak{g}$ denote by $\chi_D(t)$ the characteristic polynomial of D as a linear transformation of B_n . As is proved by Premet in [17], $\chi_D(t) = t^{p^n} + \sum_{i=0}^{n-1} \psi_i(D)t^{p^i}$ where ψ_i are algebraically independent polynomial functions generating the algebra $k[\mathfrak{g}]^{\mathcal{G}}$. There exists an open subset of $\mathfrak{g}$ consisting of elements D such that $\dim \mathfrak{g}_D = n$ and $\mathfrak{g} = \text{Lie } \mathcal{G} \oplus \mathfrak{g}_D$ where $\mathfrak{g}_D$ is the centralizer of D in $\mathfrak{g}$ ([17], Lemma 1). Hence $c_{\mathfrak{g}}(\mathfrak{g}) = d - n$ where $d = \dim \mathfrak{g}$, and $\psi_0, \dots, \psi_{n-1}$ are $\mathfrak{g}$ -invariant according to Corollary 5.3. Let $\varphi : \mathfrak{g} \rightarrow \mathbb{A}^n$ be the morphism with components $\psi_0, \dots, \psi_{n-1}$ and U the subset of those $D \in \mathfrak{g}$ for which $d_D \psi_0, \dots, d_D \psi_{n-1}$ are linearly independent. Premet proved that each fibre $F_D = \varphi^{-1}(\varphi(D))$ is irreducible of dimension $d - n$ and $F_D \cap U$ is nonempty ([17], Lemmas 12 and 13).

Put $U_1 = \{D \in \mathfrak{g} \mid \psi_0(D) \neq 0\}$. If $D \in U_1$, then D is a linear combination of $D^p, \dots, D^{p^n}$ as $D^{p^n} + \sum_{i=0}^{n-1} \psi_i(D)D^{p^i} = 0$. Hence D is semisimple, and we can find its eigenvectors $y_1, \dots, y_n \in B_n$ such that $B_n = \langle y_1, \dots, y_n \rangle \oplus (k + \mathfrak{n}^2)$. Then the monomials $y_1^{m_1} \dots y_n^{m_n}$ with $0 \leq m_i < p$ constitute a basis for B_n . Let λ_i be the eigenvalue of y_i . Since the rank of D as a linear transformation is equal to $p^n - 1$,

the equality $\sum m_i \lambda_i = 0$ can hold only for $m_1 = \dots = m_n = 0$. This implies that D generates a torus of dimension n in $\mathfrak{g}$. If $D' \in \mathfrak{g}_D$ then

$$\sum_{i=0}^{n-1} (d_D \psi_i)(D') D^{p^i} = -\psi_0(D) D',$$

which is a special case of [17], Lemma 7, (i). Taking $D' = D^{p^j}$ where $0 \leq j < n$, we see that $(d_D \psi_i)(D') \neq 0$ only for $i = j$. Hence $U_1 \subset U$. Suppose that Z is an irreducible component of the closed subset $\mathfrak{g} \setminus U_1$ having codimension 1 in $\mathfrak{g}$ (in fact it can be shown that ψ_0 is irreducible). Note that $\overline{\varphi(Z)} \neq \mathbb{A}^n$ as $\psi_0(Z) = \{0\}$. By the theorem on dimensions of fibres we have $\dim Z \cap F_D \geq \dim Z - \dim \overline{\varphi(Z)} \geq d - n$ for all $D \in Z$. It follows that Z is a union of fibres of φ . In particular, $Z \cap U \neq \emptyset$. We deduce that $\text{codim}_{\mathfrak{g}} \mathfrak{g} \setminus U \geq 2$. Thus we meet the hypotheses of Theorem 5.4:

COROLLARY 5.6. *If A denotes the algebra of polynomial functions on W_n then $A^{W_n} = A^{(p)}[\psi_0, \dots, \psi_{n-1}]$. Moreover, A^{W_n} is free of rank p^n over $A^{(p)}$ and is a locally complete intersection.*

In conclusion we make comments concerning the results of section 4. Assume that $\dim \mathfrak{g} < \infty$. According to [7] the algebra $u(\mathfrak{g}_x)$ is semisimple if and only if $\mathfrak{g}_x$ is a torus. Thus Theorem 4.2 says that $\mathfrak{g}_x$ is toral if and only if x lies in an affine open subset $U \subset X$ such that $k[U]$ is an injective $u(\mathfrak{g})$ -module. Such points x exist if and only if $k(X)$ is an injective $u(\mathfrak{g})$ -module. A. Premet pointed out to me that the openness of the set of points with a toral $\mathfrak{g}_x$ can be proved by geometric arguments.

References

1. N. Bourbaki, Commutative Algebra, Springer, Berlin, 1989.
2. E. Cline, B. Parshall and L. Scott, A Mackey imprimitivity theory for algebraic groups, Math. Z. **182** (1983) 447–471.
3. M. Demazure and P. Gabriel, Groupes Algébriques I, Masson, Paris, 1970.
4. Y. Doi, Algebras with total integrals, Comm. Algebra **13** (1985) 2137–2159.
5. S. Donkin, Infinitesimal invariants of algebraic groups, J. London Math. Soc. **45** (1992) 481–490.
6. E.M. Friedlander and B.J. Parshall, Rational actions associated to the adjoint representation, Ann. Sci. École Norm. Sup. **20** (1987) 215–226.
7. G.P. Hochschild, Representations of restricted Lie algebras of characteristic p , Proc. Amer. Math. Soc. **5** (1954) 603–605.
8. G.P. Hochschild, Simple algebras with purely inseparable splitting fields of exponent 1, Trans. Amer. Math. Soc. **79** (1955) 477–489.
9. J. Jantzen, Representations of algebraic groups, Academic Press, 1987.
10. M. Koppinen and T. Neuvonen, An imprimitivity theorem for Hopf algebras, Math. Scand. **41** (1977) 193–198.
11. H. Matsumura, Commutative Algebra, Second Edition, Benjamin, 1980.
12. A.A. Mil'ner, Irreducible representations of modular Lie algebras, Izv. Akad. Nauk SSSR Ser. Mat. **39** (1975) 1240–1259 [In Russian]. Translation in Math. USSR Izv. **9** (1975) 1169–1187.
13. S. Montgomery, Hopf algebras and Their Actions on Rings, CBMS Regional Conference Series in Mathematics, Vol. 82, American Mathematical Society, 1993.

14. D. Mumford, *Abelian Varieties*, Oxford University Press, Oxford, 1970.
15. W.D. Nichols and M.B. Zoeller, A Hopf algebra freeness theorem, *Amer. J. Math.* **111** (1989) 381–385.
16. U. Oberst and H.-J. Schneider, Über Untergruppen endlicher algebraischer Gruppen, *Manuscripta Math.* **8** (1973) 217–241.
17. A.A. Premet, The theorem on restriction of invariants and nilpotent elements in W_n , *Mat. Sbornik* **182** (1991) 746–773 [In Russian]. Translation in *Math. USSR Sbornik* **73** (1992) 135–159.
18. R. Steinberg, Regular elements of semisimple algebraic groups, *IHES Publ. Math.* **25** (1965) 49–80.
19. M.E. Sweedler, *Hopf Algebras*, Benjamin, New York, 1969.
20. M. Takeuchi, Relative Hopf modules—equivalences and freeness criteria, *J. Algebra* **60** (1979) 452–471.
21. F.D. Veldkamp, The center of the universal enveloping algebra of a Lie algebra in characteristic p , *Ann. Sci. École Norm. Sup.* **5** (1972) 217–240.
22. W.C. Waterhouse, Geometrically reductive affine group schemes, *Arch. Math.* **62** (1994) 306–307.